

# UC Davis

## UC Davis Previously Published Works

### Title

Spatially structured genetic diversity of the Amerindian yam (*Dioscorea trifida* L.) assessed by SSR and ISSR markers in Southern Brazil

### Permalink

<https://escholarship.org/uc/item/5246x01v>

### Journal

Genetic Resources and Crop Evolution, 60(8)

### ISSN

0925-9864

### Authors

Nascimento, Wellington F  
Rodrigues, Jucelene F  
Koehler, Samantha  
et al.

### Publication Date

2013-12-01

### DOI

10.1007/s10722-013-0008-y

Peer reviewed

Dear Author,

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For fax submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/ corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style. Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

#### **Please note**

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL: [http://dx.doi.org/\[DOI\]](http://dx.doi.org/[DOI]).

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information go to: <http://www.springerlink.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us if you would like to have these documents returned.

# Metadata of the article that will be visualized in OnlineFirst

---

**Please note: Images will appear in color online but will be printed in black and white.**

---

ArticleTitle                      Spatially structured genetic diversity of the endangered Amerindian yam (*Dioscorea trifida* L.) assessed by SSR and ISSR markers in Southern Brazil

---

Article Sub-Title

---

Article CopyRight                Springer Science+Business Media Dordrecht  
(This will be the copyright line in the final PDF)

---

Journal Name                      Genetic Resources and Crop Evolution

---

Corresponding Author          Family Name                      **Veasey**  
Particle  
Given Name                        **Elizabeth A.**  
Suffix  
Division                              Department of Genetics, Luiz de Queiroz College of Agriculture  
Organization                        University of São Paulo  
Address                                Av. Pádua Dias 11, CP 83, 13400-970, Piracicaba, São Paulo, Brazil  
Email                                    eaveasey@usp.br

---

Author                                Family Name                      **Nascimento**  
Particle  
Given Name                        **Wellington F.**  
Suffix  
Division                              Department of Genetics, Luiz de Queiroz College of Agriculture  
Organization                        University of São Paulo  
Address                                Av. Pádua Dias 11, CP 83, 13400-970, Piracicaba, São Paulo, Brazil  
Email

---

Author                                Family Name                      **Rodrigues**  
Particle  
Given Name                        **Jucelene F.**  
Suffix  
Division                              Department of Genetics, Luiz de Queiroz College of Agriculture  
Organization                        University of São Paulo  
Address                                Av. Pádua Dias 11, CP 83, 13400-970, Piracicaba, São Paulo, Brazil  
Email

---

Author                                Family Name                      **Koehler**  
Particle  
Given Name                        **Samantha**  
Suffix  
Division                              Department of Biological Sciences  
Organization                        Federal University of São Paulo  
Address                                Rua Professor Artur Riedel 275, 09972-270, Diadema, São Paulo, Brazil  
Email

---

Author                                Family Name                      **Gepts**  
Particle  
Given Name                        **Paul**

Suffix  
Division Department of Plant Sciences/MS1, Section of Crop and Ecosystem Sciences  
Organization University of California  
Address 1 Shields Avenue, 95616-8780, Davis, CA, USA  
Email

---

Schedule  
Received 21 February 2013  
Revised  
Accepted 22 May 2013

---

Abstract *Dioscorea trifida* L. (Dioscoreaceae) is among the economically most important cultivated Amerindian yam species, whose origin and domestication are still unresolved issues. In order to estimate the genetic diversity maintained by traditional farmers in Brazil, 53 accessions of *D. trifida* from 11 municipalities in the states of São Paulo, Santa Catarina, Mato Grosso and Amazonas were characterized based on eight Simple Sequence Repeats (SSR) and 16 Inter Simple Sequence Repeats (ISSR) markers. The level of polymorphism among the accessions was high, 95 % for SSR and 75.8 % for ISSR. The SSR marker showed higher discrimination power among accessions compared to ISSR, with *D* parameter values of 0.79 and 0.44, respectively. Although SSR and ISSR markers led to dendrograms with different topologies, both separated the accessions into three main groups: I—Ubatuba-SP; II—Iguape-SP and Santa Catarina; and III—Mato Grosso. The accessions from Amazonas State were classified in group II with SSR and in a separate group with ISSR. Bayesian and principal coordinate analyzes conducted with both molecular markers corroborated the classification into three main groups. Higher variation was found within groups in the AMOVA analysis for both markers (66.5 and 60.6 % for ISSR and SSR, respectively), and higher Shannon diversity index was found for group II with SSR. Significant but low correlations were found between genetic and geographic distances ( $r = 0.08$ ;  $p = 0.0007$  for SSR and  $r = 0.16$ ;  $p = 0.0002$  for ISSR). Therefore, results from both markers showed a slight spatially structured genetic diversity in *D. trifida* accessions maintained by small traditional farmers in Brazil.

---

Keywords (separated by '-') *Dioscorea trifida* - Genetic diversity - Genetic structure - Molecular markers - Traditional agriculture - Yams

---

Footnote Information

---

Journal: 10722  
Article: 8



## Author Query Form

**Please ensure you fill out your response to the queries raised below  
and return this form along with your corrections**

Dear Author

During the process of typesetting your article, the following queries have arisen. Please check your typeset proof carefully against the queries listed below and mark the necessary changes either directly on the proof/online grid or in the 'Author's response' area provided below

<b>Query</b>	<b>Details required</b>	<b>Author's response</b>
1.	Please check and confirm that the authors and their respective affiliations have been correctly identified and amend if necessary.	
2.	Reference Milbourne (1998) is given in list but not cited in text. Please cite in text or delete from list.	
3.	Please check the layout of Table 5, and correct if necessary.	
4.	As per the information provided by the publisher, Figs.2 and 3 will be black and white in print; hence, please confirm whether we can add "colour figure online" to the caption.	

2 **Spatially structured genetic diversity of the endangered**  
3 **Amerindian yam (*Dioscorea trifida* L.) assessed by SSR**  
4 **and ISSR markers in Southern Brazil**

5 **Wellington F. Nascimento · Jucelene**  
6 **F. Rodrigues · Samantha Koehler · Paul Gepts ·**  
7 **Elizabeth A. Veasey**

8 Received: 21 February 2013 / Accepted: 22 May 2013  
9 © Springer Science+Business Media Dordrecht 2013

10 **Abstract** *Dioscorea trifida* L. (Dioscoreaceae) is  
11 among the economically most important cultivated  
12 Amerindian yam species, whose origin and domesti-  
13 cation are still unresolved issues. In order to estimate  
14 the genetic diversity maintained by traditional farmers  
15 in Brazil, 53 accessions of *D. trifida* from 11 munic-  
16 ipalities in the states of São Paulo, Santa Catarina,  
17 Mato Grosso and Amazonas were characterized based  
18 on eight Simple Sequence Repeats (SSR) and 16 Inter  
19 Simple Sequence Repeats (ISSR) markers. The level of  
20 polymorphism among the accessions was high, 95 %  
21 for SSR and 75.8 % for ISSR. The SSR marker showed  
22 higher discrimination power among accessions com-  
23 pared to ISSR, with *D* parameter values of 0.79 and  
24 0.44, respectively. Although SSR and ISSR markers  
25 led to dendrograms with different topologies, both  
26 separated the accessions into three main groups:

I—Ubatuba-SP; II—Iguape-SP and Santa Catarina; 27  
and III—Mato Grosso. The accessions from Amazonas 28  
State were classified in group II with SSR and in a 29  
separate group with ISSR. Bayesian and principal 30  
coordinate analyzes conducted with both molecular 31  
markers corroborated the classification into three main 32  
groups. Higher variation was found within groups in 33  
the AMOVA analysis for both markers (66.5 and 34  
60.6 % for ISSR and SSR, respectively), and higher 35  
Shannon diversity index was found for group II with 36  
SSR. Significant but low correlations were found 37  
between genetic and geographic distances ( $r = 0.08$ ; 38  
 $p = 0.0007$  for SSR and  $r = 0.16$ ;  $p = 0.0002$  for 39  
ISSR). Therefore, results from both markers showed a 40  
slight spatially structured genetic diversity in *D. trifida* 41  
accessions maintained by small traditional farmers in 42  
Brazil. 43

A1 W. F. Nascimento · J. F. Rodrigues · E. A. Veasey (✉)  
A2 Department of Genetics, Luiz de Queiroz College  
A3 of Agriculture, University of São Paulo, Av. Pádua Dias  
A4 11, CP 83, Piracicaba, São Paulo 13400-970, Brazil  
A5 e-mail: eaveasey@usp.br

A6 S. Koehler  
A7 Department of Biological Sciences, Federal University of  
A8 São Paulo, Rua Professor Artur Riedel 275, Diadema, São  
A9 Paulo 09972-270, Brazil

A10 P. Gepts  
A11 Department of Plant Sciences/MS1, Section of Crop  
A12 and Ecosystem Sciences, University of California,  
A13 1 Shields Avenue, Davis, CA 95616-8780, USA

**Keywords** *Dioscorea trifida* · Genetic diversity · 44  
Genetic structure · Molecular markers · Traditional 45  
agriculture · Yams 46

**Introduction** 49

The genus *Dioscorea*, family Dioscoreaceae, repre- 50  
sents an important food source in the humid and 51  
subhumid tropics (Ayensu and Coursey 1972). This 52  
genus consists of more than 600 species, of which only 53  
10 are used for human consumption (Lebot 2009). 54  
*Dioscorea trifida* L., originating in South America, is 55

56 among the economically most important species,  
57 including *Dioscorea cayenensis* Lam. and *Dioscorea*  
58 *rotundata* Poir., originating in Africa, and *Dioscorea*  
59 *alata* L., originating in Asia (Coursey 1976).

60 *Dioscorea trifida* is an herbaceous, autotetraploid  
61 ( $x = 20$  and  $2n = 4x = 80$ ), viny, and perennial plant  
62 (Bousalem et al. 2006), with quadrangular winged  
63 stems without spines but with deeply lobed leaves,  
64 usually arranged alternately or rarely opposite (Mont-  
65 aldo 1991). The plants are dioecious, with small  
66 unisexual flowers that, when fertilized, produce ined-  
67 ible encapsulated fruits (Stephens 2009). Its repro-  
68 duction occurs by allogamy or vegetative propagation  
69 (Montaldo 1991). The tuber, the edible plant structure,  
70 has a high nutritional quality and astringent, antimicro-  
71 bial and diuretic properties, which allow its use for  
72 combating malnutrition and treatment of diseases such  
73 as diabetes and high cholesterol levels (Ramos-  
74 Escudero et al. 2010). The main limiting factor for  
75 growing *D. trifida* is potyviruses (genus *Potyvirus*,  
76 family Potviridae), which causes a variety of symp-  
77 toms on the leaves of infected plants (Odu et al. 2004).  
78 Potyvirus infection can cause significant economic  
79 damage and process of genetic erosion of the crop  
80 (Bousalem et al. 2010).

81 The evolutionary history of *D. trifida* is controver-  
82 sial. Although it occurs very frequently in various  
83 countries of Latin America, and the Amazon has been  
84 reported as a possible center of origin and diversifi-  
85 cation of this species (Degras 1993), the lack of  
86 information about its origin and domestication process  
87 is still evident. It is believed that *D. trifida* originated  
88 on the border between Brazil, Guyana, French Guyana  
89 and Suriname, and was domesticated by indigenous  
90 peoples in these regions (Pedralli 1998). Recent  
91 studies conducted in French Guiana revealed the  
92 presence of wild relatives of *D. trifida*, being the first  
93 direct genetic evidence of possible places of origin for  
94 this species (Bousalem et al. 2010).

95 In the Amazon, Clement (1999) observed the  
96 existence of several areas with large concentrations  
97 of genetic resources related to different crop species.  
98 *D. trifida* was present in some of these sites, such as in  
99 the Northwestern Amazonian Center, the Central  
100 Amazonian Center, the Middle Orinoco Minor Centre  
101 and the Guiana Minor Centre, indicating the close  
102 relationship of these areas with the evolutionary  
103 history of *D. trifida*. In archaeological excavations in  
104 Panama, *D. trifida* tubers were found together with

105 cassava (Piperno et al. 2000, Dickau et al. 2007). As  
106 cassava was domesticated in southwestern Amazon  
107 Basin (Olsen 2004) and quickly spread throughout  
108 Tropical America (Piperno et al. 2000, Dickau et al.  
109 2007), *D. trifida* could have been domesticated and  
110 propagated by the same tribes involved in the process  
111 of cassava domestication (Bousalem et al. 2010),  
112 possibly being the first yam species cultivated by  
113 indigenous peoples in the Amazon (Degras 1993). The  
114 cultivation of *D. trifida* in Brazil has taken place since  
115 then, mainly by small rural farmers (Pedralli 1998). In  
116 recent surveys, the occurrence of this species was  
117 observed in the Central West, South and Southeast  
118 Brazil (Bressan et al. 2005; Veasey et al. 2010). It is  
119 also an important crop in the Amazon (Velez 1998).

120 Despite the geo-cultural and socioeconomic impor-  
121 tance of *D. trifida*, few studies are conducted to  
122 explore its potential and to develop conservation  
123 strategies for this crop. In Brazil, there are few  
124 institutions currently involved in research related to  
125 the yam crop; therefore, new studies are important to  
126 add information for breeding programs and conserva-  
127 tion strategies. As the cultivation and consumption of  
128 yam are very intense in family agricultural production  
129 systems practiced by traditional communities, these  
130 systems provide a favorable environment for the  
131 generation and maintenance of genetic diversity of this  
132 crop (Veasey et al. 2010). However, the socioeco-  
133 nomic pressures faced by farmers in recent years have  
134 caused the loss of plant genetic resources, specifically  
135 *D. trifida*, and biodiversity losses can be severe and  
136 irreversible. In this context, there is a need to estimate  
137 the genetic diversity of *D. trifida* maintained by  
138 traditional farmers to assist in developing strategies to  
139 preserve the species and lessen losses caused by  
140 various socioeconomic pressures on the yam crop.

141 Various molecular biological techniques are avail-  
142 able to detect genetic variability of natural populations  
143 and cultivated plants. Among these techniques, micro-  
144 satellites or Simple Sequence Repeats (SSR) are very  
145 effective because they are codominant, multi-allelic,  
146 highly polymorphic and show good reproducibility  
147 (Oliveira et al. 2006). In order to study the genetic  
148 diversity of *D. trifida*, Hochu et al. (2006) developed  
149 eight SSR primers specific for this species, which were  
150 used for the analysis of 24 cultivars, showing high  
151 polymorphism. These primers were used by Bousalem  
152 et al. (2006) to assess the inheritance pattern of  
153 *D. trifida*, from the analysis of parental genotypes and



154 offspring, where the tetraploid behavior of the species  
155 was reported.

156 Another marker used in genetic diversity studies is  
157 Inter Simple Sequence Repeats (ISSR), which were  
158 developed to explore microsatellite repeats without  
159 the need to use of DNA sequencing (Zietkiewicz et al.  
160 1994; Reddy et al. 2002). ISSR markers are very  
161 stable, dominant, multi-allelic, present good repro-  
162 ducibility and generate a large number of polymorphic  
163 fragments (Mattioni et al. 2002; Wolfe 2005). Few  
164 genetic diversity studies have been performed in the  
165 *Dioscorea* genus based on ISSR molecular markers.  
166 Among them, Zhou et al. (2008) analyzed the level of  
167 genetic diversity among different cultivars of *Diosco-*  
168 *rea opposita* Thunb., widely used in traditional  
169 Chinese medicine and Wu et al. (2009) evaluated the  
170 relationship and genetic variability among accessions  
171 of *D. alata*. Both studies found that ISSR provided a  
172 good assessment of genetic diversity of yam and  
173 valuable information to help in selecting parents for  
174 future yam breeding programs.

175 The aim of this study was to characterize 53  
176 accessions of *D. trifida* originating in traditional  
177 communities in the States of Santa Catarina, São  
178 Paulo, Mato Grosso and Amazonas, using ISSR and  
179 SSR markers, in order to verify the level of genetic  
180 diversity maintained by farmers in these regions in  
181 Brazil. The study describes the spatially structured  
182 genetic variation of *D. trifida* maintained by these  
183 farmers and the genetic diversity that is concentrated  
184 within the different sampling sites.

## 185 Materials and methods

### 186 Plant materials

187 We evaluated 53 accessions of *D. trifida* collected  
188 from 11 municipalities in the States of Sao Paulo (SP),  
189 Santa Catarina (SC), Mato Grosso (MT) and Amazo-  
190 nas (AM), located between latitudes 14°43'S and  
191 26°15'S and longitudes 44°01'W and 62°05'W (Fig. 1;  
192 Table 1). In each visited municipality, a collection  
193 was conducted so as to seek greater representation of  
194 the genetic variability, taking into account morpho-  
195 logical variation and information from farmers. Three  
196 accessions (the two accessions from Amazonas and  
197 one from Ubatuba, SP) were acquired in local markets.  
198 Accessions were collected in the form of tubers, which

were grown in pots placed in a greenhouse at the Luiz  
de Queiroz College of Agriculture, University of Sao  
Paulo, in Piracicaba, SP, located at 22°43'S latitude  
and 47°25'W longitude, where young leaves were  
collected for DNA extraction.

### DNA extraction and quantification

Young, newly expanded leaves were collected and  
stored at 4 °C for 7 days in a CTAB gel, containing  
30 mg CTAB, 350 mg NaCl and 70 ml of distilled  
water (Rogstad 1992). After this period, the gel excess  
was removed from plant tissues with the aid of a paper  
towel. The fragments were then macerated in 1 mL  
STE buffer [0.13 mg saccharose, 45 mL of Tris-HCL  
(1 M), 150 mL of EDTA (0.5 M), completing with  
distilled water to a final volume of 1.5 mL] and  
subjected to DNA extraction by the method of Doyle  
and Doyle (1990). DNA concentration was estimated  
in a 1 % agarose gel, using a TBE 10X running buffer,  
stained in ethidium bromide. A final concentration of  
5 ng/μL was obtained for the PCR analysis.


### Amplification of SSR and ISSR

For the SSR amplification, 10 primer pairs developed  
by Hochu et al. (2006) and Tostain et al. (2006) were  
tested (Table 2). PCR was conducted in a 16 μL  
reaction volume containing: 5 ng of genomic DNA in  
a 5× reaction buffer, 1.5 mM MgCl<sub>2</sub>, 2.5 mM dNTPs,  
5 pmol of forward primer, 5 pmol of reverse primer,  
and 5 U/μL *Taq* DNA polymerase (Promega, Madi-  
son, USA). The amplification reactions were per-  
formed in a MyCycler Thermal Cycler model BioRad  
thermocycler using the following steps: 1) denatur-  
ation at 94 °C for 5 min, followed by 30 cycles [30 s  
at 94 °C, 30 s at annealing temperature (touchdown of  
50–60 °C) and 30 s for 72 °C], and a final stage of  
extension of 5 min at 72 °C for the PCR reactions with  
specific primers for *D. trifida* (Hochu et al. 2006); 2)  
denaturation at 94 °C for 5 min, followed by 35 cycles  
[30 s at 94 °C, 1 min at a temperature of annealing  
(touchdown 50–60 °C) and 1 min at 72 °C], and a  
final extension of 8 min at 72 °C for PCR reactions  
with heterologous primers (Tostain et al. 2006).

Electrophoresis was performed on denaturing 7 %  
polyacrylamide gel, with a constant power of 70 W for  
the time necessary for separating the amplified frag-  
ments in each primer, using 10 and 100 bp DNA



**Fig. 1** Collection sites of *D. trifida* in Brazil. Details of each accession are given in Table 1



244 Ladder (Invitrogen™, São Paulo, Brazil) as markers.  
 245 The gels were stained using silver nitrate methodology  
 246 (Creste et al. 2001) for the revelation of microsatellite  
 247 bands, which were photographed with a digital camera  
 248 and evaluated in a transilluminator.

249 For the ISSR analysis 20 primers were tested  
 250 according to the Wolfe (2000, 2005) protocol  
 251 (Table 3). PCR reactions were performed in a final  
 252 volume of 30 µL containing: 5 ng of genomic DNA in  
 253 a 5× reaction buffer, 2.0 mM MgCl<sub>2</sub>, 2.33 mM  
 254 dNTPs, 10 pmol primer and 5 U/µL Taq DNA  
 255 polymerase (Promega, Madison, USA). The amplifi-  
 256 cation of the DNA template were performed in  
 257 MultiGene Thermal Cycler thermocycler (Labnet  
 258 International, Inc.) according to the following ampli-  
 259 fication conditions: 90 s at 94 °C, 35 cycles at 94 °C  
 260 for 40 s, followed by 46 cycles (52 °C for 45 s, 72 °C  
 261 for 90 s, 94 °C for 45 s, 44 °C for 45 s), and a final  
 262 stage of extension at 72 °C for 5 min (Wolfe 2000).

263 The products resulting from the amplification  
 264 reactions were subjected to electrophoresis on a 2 %  
 265 agarose gel in TBE buffer 10X for 140 min at 90 V

266 and stained with ethidium bromide. A 100 bp DNA  
 267 Ladder (Invitrogen™, Carlsbad, USA) was used as a  
 268 marker. Additionally, we used control samples previ-  
 269 ously amplified with success. The gel was photo-  
 270 graphed over ultraviolet light source with Syngene  
 271 photodocumentation system (Synoptics Ltda., Cam-  
 272 bridge, United Kingdom). For statistical analysis we  
 273 considered only robust and unambiguous bands. We  
 274 discarded the bands that showed low intensity or  
 275 coalescing with other bands.

#### 276 Statistical analysis 276

277 Due to the tetraploid behavior of *D. trifida*, as  
 278 described by Bousalem et al. (2006), the band patterns  
 279 of the SSR and ISSR markers were both interpreted as  
 280 binary data, presence (1) and absence (0) of bands,  
 281 generating data matrices that were subjected to the  
 282 following statistical programs. Genetic diversity anal-  
 283 yses were based on POPGENE Software, version 1.3  
 284 (Yeh et al. 1997), where we obtained the number of  
 285 bands observed per primer, number of polymorphic

**Table 1** *Dioscorea trifida* accessions used in this study collected in Brazil, including accession and identification number (ID) in the Germplasm Bank, origin (community, municipality, state), popular name and geographic coordinates

Accession	ID	Geographic localities	Popular name	Lat/Long
01	180	Sertão de Ubatumirim, Ubatuba, São Paulo	Cará roxo	23°15'S/44°01'O
02	181	Sertão de Ubatumirim, Ubatuba, São Paulo	Cará branco	23°17'S/44°05'O
03	182	Sertão de Ubatumirim, Ubatuba, São Paulo	Cará roxo	23°18'S/44°52'O
04	183	Sertão de Ubatumirim, Ubatuba, São Paulo	Cará roxo	23°17'S/44°51'O
05	184	Sertão de Ubatumirim, Ubatuba, São Paulo	Cará roxo	23°18'S/44°51'O
06	185	Sertão de Ubatumirim, Ubatuba, São Paulo	Cará roxo	23°18'S/44°51'O
07	187	Sertão de Ubatumirim, Ubatuba, São Paulo	Cará branco	23°18'S/44°51'O
08	191	Sertão das Cutias, Ubatuba, São Paulo	Cará roxo	23°22'S/44°58'O
09	193	Rio Escuro, Ubatuba, São Paulo	Cará branco	23°28'S/45°08'O
10	195	Sertão do Ingá, Ubatuba, São Paulo	Cará cobrinha	23°31'S/45°13'O
11	196	Sertão do Ingá, Ubatuba, São Paulo	Cará branco	23°31'S/45°13'O
12	197	Sertão de Ubatumirim, Ubatuba, São Paulo	Cará roxo	23°17'S/44°51'O
13	198	Sertão do Ingá, Ubatuba, São Paulo	Cará roxo	23°31'S/45°13'O
14	201	Sertão do Ingá, Ubatuba, São Paulo	Cará roxo	23°31'S/45°14'O
15	203	Sertão do Ingá, Ubatuba, São Paulo	Cará roxo	23°31'S/45°14'O
16	204	Rio Escuro, Ubatuba, São Paulo	Cará roxo	23°28'S/45°08'O
17	208	Araribá, Ubatuba, São Paulo	Cará roxo	23°32'S/45°15'O
18	210	Sertão de Ubatumirim, Ubatuba, São Paulo	Cará roxo	23°29'S/45°10'O
19	216	Fazenda da Caixa, Ubatuba, São Paulo	Cará roxo	23°31'S/45°14'O
20	217	Feira de Ubatuba, Ubatuba, São Paulo	Cará roxo	23°27'S/45°09'O
21	236	Feira de Manaus, Manaus, Amazonas	Cará roxo	03°08'S/60°01'O
22	237	Feira de Barcelos, Barcelos, Amazonas	Cará	0°58'S/62°55'O
23	281	Pirabeiraba, Joinville, Santa Catarina	Cará	26°10'S/48°55'O
24	282	Pirabeiraba, Joinville, Santa Catarina	Cará mimoso	26°09'S/48°56'O
25	283	Pirabeiraba, Joinville, Santa Catarina	Cará	26°09'S/48°58'O
26	285	Acaraí, São Francisco do Sul, Santa Catarina	Cará pão	26°11'S/48°53'O
27	286	Pirabeiraba, Joinville, Santa Catarina	Cará mimoso	26°15'S/48°37'O
28	287	Pirabeiraba, Joinville, Santa Catarina	Carcanhá de nego	26°09'S/48°59'O
29	290	Pirabeiraba, Joinville, Santa Catarina	Cará mimoso	26°09'S/48°59'O
30	292	Pirabeiraba, Joinville, Santa Catarina	Cará	26°09'S/48°59'O
31	297	Pirabeiraba, Joinville, Santa Catarina	Cará	26°09'S/48°59'O
32	298	Rio da Prata, Joinville, Santa Catarina	Cará	26°11'S/48°58'O
33	301	Rio da Prata, Joinville, Santa Catarina	Cará mimoso	26°11'S/48°58'O
34	302	Pirabeiraba, Joinville, Santa Catarina	Cará	26°10'S/48°57'O
35	312	Icapara, Iguape, São Paulo	Cará São João branco	24°40'S/47°27'O
36	313	Cavalcanti, Iguape, São Paulo	Cará-pipa	24°43'S/47°45'O
37	323	Icapara, Iguape, São Paulo	Cará São João roxo	24°40'S/47°27'O
38	328	Momuna, Iguape, São Paulo	Cará São João roxo	24°42'S/47°40'O
39	329	Momuna, Iguape, São Paulo	Cará São João branco	24°42'S/48°40'O
40	335	Carumbé, Acorizal, Mato Grosso	Cará roxo	15°08'S/56°12'O
41	336	Carumbé, Acorizal, Mato Grosso	Cará roxo	15°08'S/56°12'O
42	340	Rio dos Couros, Cuiabá, Mato Grosso	Cará pé de anta	15°36'S/55°48'O
43	343	Carumbé, Acorizal, Mato Grosso	Cará branco	15°08'S/56°12'O

**Table 1** continued

Accession	ID	Geographic localities	Popular name	Lat/Long
44	344	Sela Dourada, Nobres, Mato Grosso	Cará do Joaquim	15°36'S/56°48'O
45	345	Santo Antônio do Barreiro, Jangada, Mato Grosso	Cará roxo	15°08'S/56°17'O
46	350	Sela Dourada, Nobres, Mato Grosso	Cará branco	15°34'S/56°46'O
47	351	Sela Dourada, Nobres, Mato Grosso	Cará mão de anta	15°30'S/56°42'O
48	352	Timbozal, Rosário Oeste, Mato Grosso	Cará mão de anta	14°51'S/56°23'O
49	355	Chapada Vacaria, Acorizal, Mato Grosso	Cará roxo	15°03'S/56°08'O
50	361	Sela Dourada, Nobres, Mato Grosso	Cará roxo	14°43'S/56°15'O
51	364	Barranco Alto, Rosário Oeste, Mato Grosso	Pombinho branco	15°14'S/57°59'O
52	366	Sela Dourada, Nobres, Mato Grosso	Cará roxo cumprido	14°43'S/56°15'O
53	368	Barranco Alto, Rosário Oeste, Mato Grosso	Cará roxo	15°17'S/57°50'O

**Table 2** List of SSR primers used to evaluate 53 *Dioscorea trifida* accessions, including primer sequence, annealing temperature ( $T_A$ ); size range of SSR bands in base pairs (bp), number of bands ( $N_B$ ), number of polymorphic bands ( $N_{PB}$ ), percent polymorphism ( $P$ ) and discriminating power ( $D$ )

Primer code	Sequence (5'-3')	$T_A$ (°C)	Size range (bp)	$N_B$	$N_{PB}$	$P$ (%)	$D$
Da1A01 <sup>1</sup>	F: TAT AAT CGG CCA GAG G R: TGT TGG AAG CAT AGA GAA	51–53	202–205	2	2	100.0	0.97
Dab2C05 <sup>1</sup>	F: CCC ATG CTT GTA GTT GT R: TGC TCA CCT CTT TAC TTG	51–52	168–192	5	5	100.0	0.91
MTI2 <sup>2</sup>	F: TCATCAAGAGCATCAAAAAAC R: GCCTCGTCTTTGAAGTTGGT	50–52	121–131	6	6	100.0	0.71
MTI3 <sup>2</sup>	F: TAACAAACAAAAATGAAAC R: TAACAGTGATTGAGCTAGGA	55–59	156–205	13	13	100.0	0.85
MTI4 <sup>2</sup>	F: ACTTGGTGTGTTGGATTGC R: TATCACTCCCCAGACCAGA	50–58	101–111	8	8	100.0	0.61
MTI10 <sup>2</sup>	F: TCGTGTCATCTTGCTGCGT R: GAAAAGCGGAGATGAAGAGCA	55–58	143–198	11	11	100.0	0.61
MTI11 <sup>2</sup>	F: CTCTTTTGCTTCTCATTTCA R: ATGTAGCCAATCCAAAATAG	55–56	124–137	5	4	80.0	0.72
MTI12 <sup>2</sup>	F: CTGCCAGCGTCCGATTC R: CGTAGGACCTCTCGCATCAG	55–60	100–123	6	5	83.0	0.92
Average		–	–	7.0	6.75	95.0	0.79

<sup>1</sup> Tostain et al. (2006); <sup>2</sup> Hochu et al. (2006)

bands, percent polymorphism and estimated the Shannon index according to the following formula:  $H' = -\sum_{i=1}^s pi \log pi$ , where  $pi$  is the frequency of each species, for  $i$  ranging from 1 to  $S$  (richness).

In order to compare the efficiency of the markers in the genotypic identification, the discrimination power ( $D$ ) (Tessier et al. 1999) was estimated for each primer.

This parameter was calculated according to the formula:  $D_j = 1 - C_j = 1 - \sum_{i=1}^I pi \frac{(Npi-1)}{N-1}$ , where  $D$  is the probability that two randomly selected individuals have a different and distinct banding pattern from each other;  $C$  is the probability that two randomly selected individuals have a similar band pattern, and  $N$  is the number of individuals analyzed.

**Table 3** ISSR primers used to evaluate 53 *Dioscorea trifida* accessions, including primer sequence, annealing temperature ( $T_A$ ), size range of ISSR bands in base pairs (bp), number of bands ( $N_B$ ), number of polymorphic bands ( $N_{PB}$ ), percent polymorphism ( $P$ ), and discriminating power ( $D$ )

Primer code	Sequence (5'-3')	$T_A$ (°C)	Size range(bp)	$N_B$	$N_{PB}$	$P$ (%)	$D$
UBC 7	(CT)8-RG	48	300–1,300	13	11	84.6	0.37
UBC 814	(CT)8-TG	50	500–1,100	8	6	75.0	0.76
UBC 843	(CT)8-RA	48	600–1,200	6	6	100.0	0.82
UBC 844	(CT)8-RC	50	300–1,300	9	2	22.2	-0.35
UBC 898	(CA)6-RY	48	300–1,300	12	6	50.0	-0.48
UBC 899	(CA)6-RG	54	300–1,300	9	7	77.8	0.63
JOHN	(AG)7-YC	54	100–1,200	13	9	69.2	0.81
UBC 901	(GT)6-YR	50	300–1,300	8	8	100.0	0.67
UBC 902	(GT)6-AY	50	500–900	4	4	100.0	0.77
AW3	(GT)6-RG	54	500–800	4	3	75.0	0.75
OMAR	(GAG)4-RC	50	300–900	8	8	100.0	0.54
DAT	(GA)7-RG	54	300–800	9	2	22.2	-0.16
TERRY	(GTG)4-RC	50	300–800	9	5	55.6	0.49
MAO	(CTC)4-RC	50	400–1,300	8	8	100.0	0.36
MANNY	(CAC)4-RC	48	300–1,000	11	9	81.8	0.48
GOOFY	(GT)7-YG	54	300–900	6	6	100.0	0.56
Average	–	–	–	8.56	6.25	75.8	0.44

300 DARwin software, version 5.0 (Perrier and Jac-  
 301 quemoud-Collet 2006), was used to perform a cluster  
 302 analysis, based on Jaccard similarity coefficient and  
 303 the UPGMA method. The stability of the groupings  
 304 was assessed based on estimates of genetic dissimi-  
 305 larity through the procedure of resampling with 1,000  
 306 bootstraps. Values higher than 70 % in the nodes that  
 307 join the groups indicate homogeneity among acces-  
 308 sions. Software NTSYS-pc (Rohlf 1992) was used to  
 309 conduct a principal coordinate analysis (PCoA) and  
 310 obtain scatter plots.

311 To confirm the reliability of the groups obtained in  
 312 the cluster analysis and PCoA, we conducted a  
 313 Bayesian analysis using the software Structure (Prit-  
 314 chard et al. 2000; Pritchard and Donnelly 2001; Falush  
 315 et al. 2007), which does not rely on prior information  
 316 on possible groups, for example based on the origin of  
 317 the accessions. The Structure software was run using  
 318 the admixture model, correlated allele frequencies and  
 319 repeated ten times for each K (number of assumed  
 320 clusters) with a burn-in of 500,000 interactions  
 321 followed by 500,000 interactions MCMC (Markov  
 322 Chain Monte Carlo). The most likely number of  
 323 clusters was chosen using the  $\Delta K$  method (Evanno  
 324 et al. 2005).

325 In order to identify the proportion of genetic  
 326 variation between and within groups obtained using  
 327 the software Structure, which coincided with the

328 groups of PCoA and cluster analysis, a molecular  
 329 variance analysis (AMOVA) was carried out with  
 330 Arlequin software (Schneider et al. 2000). Another  
 331 parameter analyzed was the correlation between  
 332 matrices of genetic and geographic distances, as well  
 333 as between genetic distance matrices for SSR and  
 334 ISSR markers, through the Pearson correlation ( $r$ ),  
 335 whose significance was evaluated by Mantel (1967)  
 336 test, using NTSYS-pc software (Rohlf 1992).

## 337 Results

338 Eight SSR and 16 ISSR primers were selected based  
 339 on the presence of well defined and with good  
 340 resolution bands (Tables 2, 3). We obtained 56 bands  
 341 or amplification products with sizes ranging from  
 342 101 pb to 205 pb for SSR and 137 bands ranging from  
 343 100 bp to 1,300 pb for ISSR, in a total of 193 bands,  
 344 with an average of 7.0 bands/primer for SSR and 8.56  
 345 bands/primer for ISSR. The number of polymorphic  
 346 bands for SSR and ISSR was 54 and 100, with an  
 347 average of 6.75 and 6.25 polymorphic bands per  
 348 primer, respectively. The level of polymorphism was  
 349 high, 95 % for SSR and 75.8 % for ISSR. Parameter  
 350  $D$  value for SSR was 0.79, while for ISSR was 0.44,  
 351 demonstrating that although the ISSR marker has  
 352 generated a greater number of bands, the SSR marker

353 showed greater discriminatory power between the  
354 accessions.


355 The Jaccard coefficient among 53 accessions of *D.*  
356 *trifida* ranged from 0.40 to 0.96, with a variation of  
357 56 % similarity for SSR marker and from 0.66 to 0.97,  
358 with a variation of 31 % for ISSR. Although the two  
359 types of markers are located mostly in neutral regions  
360 and related to different sequences of the genome, the  
361 correlation between genetic matrices obtained from  
362 SSR and ISSR markers was high ( $r = 0.57$ ;  $p =$   
363 0.0002), demonstrating similar relationships between  
364 data from both marker classes.

365 Although ISSR and SSR markers generated dendro-  
366 grams with different topologies (Figs. 2, 3), in general,  
367 both dendrograms showed the formation of the same  
368 groups, with a few exceptions. Despite the low  
369 bootstrap values, below 60 % and thus not shown in  
370 the dendrograms, it was possible to identify three well-


371 defined groups: group I (accessions from Ubatuba-  
372 SP), group II (accessions from Iguape-SP and Santa  
373 Catarina-SC) and group III (accessions from Mato  
374 Grosso-MT). The yam varieties collected in Iguape-  
375 SP and Santa Catarina showed higher genetic similar-  
376 ity, while the varieties from Ubatuba-SP and Mato  
377 Grosso were more divergent and classified into distinct  
378 groups. All accessions were grouped according to their  
379 collection locations for both markers, except the  
380 accessions from Amazonas, which changed their  
381 position in the dendrogram according to the molecular  
382 marker analyzed. These two accessions were classified  
383 into a separate group (group IV) in the ISSR analysis  
384 while in the SSR analysis they were classified in group  
385 II. Also, within group II, accessions from Santa  
386 Catarina were apparently better separated from those  
387 from Iguape-SP in the SSR than in the ISSR cluster  
388 analysis. Although variations obtained in the PCoA,

**Fig. 2** UPGMA

dendrogram based on eight  
SSR primers showing the  
genetic relationships among  
53 accessions of *D. trifida*:  
group I [accessions from  
Ubatuba-SP (*green*)]; group  
II [accessions from Iguape-  
SP (*blue*), Santa Catarina  
(*pink*) and Amazonas  
(*yellow*)]; and group III  
[accessions from Mato  
Grosso (*red*)]




**Fig. 3** UPGMA dendrogram based on 16 ISSR primers showing the genetic relationships among 53 accessions of *D. trifida*: group I [accessions from Ubatuba-SP (*green*)]; group II [accessions from Iguape-SP (*blue*) and Santa Catarina (*pink*)]; group III [accessions from Mato Grosso (*red*)] and group IV [accessions from Amazonas (*yellow*)]



389 whose first two principal coordinates represented  
390 35.7 % of total variation, were not significant, a  
391 scatter plot from data obtained with SSR separated the  
392 genotypes in the same groups obtained in the scatter  
393 plot of the data obtained from ISSR, whose first two  
394 principal coordinates represented 31.6 % (not shown  
395 data). So, both markers seem to be useful in discrim-  
396 inating the genetic diversity of *D. trifida* accessions.

397 The Bayesian analysis performed with Structure  
398 software for the SSR and ISSR data confirmed the  
399 groups obtained in the SSR cluster analysis and the  
400 PCoA, since the value of  $k$  was equal to three, showing  
401 that the accessions are genetically structured in three  
402 groups (Fig. 4). Based on SSR data, the two Amazon-  
403 ian accessions showed more than 90 % of their

404 genetic constitution similar to accessions from Iguape-  
405 SP and Santa Catarina (Fig. 4a), while based on ISSR  
406 data, the same accessions showed more than 60 %  
407 similarity to those from Mato Grosso and more than  
408 30 % to those from Ubatuba-SP (Fig. 4b). Comparing  
409 both analysis, the SSR marker (Fig. 4a) showed some  
410 exceptions to the groups formed in the dendrogram for  
411 this marker (Fig. 2), such as accessions no. 17 and 18,  
412 from Ubatuba-SP, showing more than 50 % similarity  
413 to the group from Iguape-SP and Santa Catarina, while  
414 accession no. 40, from Mato Grosso, showed more  
415 than 80 % similarity to the group from Iguape-SP and  
416 Santa Catarina. The ISSR marker (Fig. 4b), on the  
417 other hand, showed a group pattern with great  
418 similarity to the ISSR dendrogram in Fig. 3.



**Fig. 4** Structure of the genetic diversity of 53 *D. trifida* accessions based on Bayesian approach for eight SSR markers (a) and 16 ISSR markers (b). Each accession is represented by a

vertical bar. Accessions 1–20 originated from Ubatuba-SP; accessions 21–39 originated from Iguape-SP, Santa Catarina and Amazonas; and accessions 40–53 originated from Mato Grosso

419 From the analysis of variance considering the three  
420 groups formed in the Bayesian analysis and two  
421 comparison levels, within and between groups, dif-  
422 ferences were found between the genetic material  
423 studied for both the SSR data ( $\Phi_{st} = 0.39$ ;  $p =$   
424  $0.0000$ ) and for ISSR data ( $\Phi_{st} = 0.33$ ;  $p = 0.0000$ )  
425 (Table 4). Genetic variation was greater within than  
426 between groups for both SSR (60.6 %) and ISSR  
427 (66.5 %) markers (Table 4). A low positive correla-  
428 tion was identified between genetic and geographical  
429 distances for both SSR data ( $r = 0.08$ ;  $p = 0.0007$ )  
430 and ISSR data ( $r = 0.16$ ;  $p = 0.0002$ ), demonstrat-  
431 ing a slight spatial structure of genetic material in the  
432 geographic area sampled.

433 The Shannon diversity index obtained from SSR  
434 data indicated that the State of São Paulo showed  
435 greater genetic diversity (0.40), followed by Mato  
436 Grosso (0.31) and Santa Catarina (0.27) (Table 5).  
437 Similar results were obtained for ISSR data, but with  
438 lower values (0.28, 0.21 and 0.19, respectively). The  
439 Amazonas State had the lowest diversity indexes (0.09  
440 for SSR and 0.03 for ISSR), due to the low number of  
441 accessions sampled. Considering the groups formed in  
442 the Bayesian analysis and in the cluster analysis with  
443 SSR, group II (accessions from Iguape-SP, Santa  
444 Catarina and Amazonas) had the highest diversity  
445 index (0.40), followed by group III (0.31) with acces-  
446 sions from Mato Grosso, and group I (0.30) (accessions  
447 from Ubatuba-SP) for SSR, while lower and similar  
448 values for the three groups were obtained for ISSR data.

**Table 4** Analysis of molecular variance (AMOVA) for 53 accessions of *D. trifida* L. considering three groups, according to the Bayesian analysis on Structure software: group I (accessions from Ubatuba-SP), group II (accessions from Iguape-SP, Santa Catarina and Amazonas), and group III (accessions from Mato Grosso)

Variation source	DF	SSR		ISSR	
		SQ	Total variation (%)	SQ	Total variation (%)
Among groups	3	150.295	39.37	221.218	33.53
Within group	49	288.346	60.63	528.404	66.47
Total	52	438.642			

$\Phi_{st} = 0.3937$  for SSR and  $\Phi_{st} = 0.3353$  for ISSR; *DF* degrees of freedom, *SQ* sum of squares

\* Value  $p^1(1,023$  permutations) = 0.0000

## Discussion

Genetic diversity and comparative analysis of SSR and ISSR markers

The large amount of molecular markers available to estimate genetic diversity allows us to make comparisons in order to determine which technique is best suited for a particular crop (Biswas et al. 2010). Choosing the most appropriate technique depends on the purpose of the research, reproduction mode of the

**Table 5** Shannon diversity index ( $H'$ ) of accessions of *D. trifida* classified according to State, to both municipalities in São Paulo (Ubatuba and Iguape), and to Bayesian analysis on Structure software: group I (accessions from Ubatuba-SP), group II (accessions from Iguape-SP, Santa Catarina and Amazonas), and group III (accessions from Mato Grosso)

	$H'$	
	SSR	ISSR
States		
São Paulo	0.40	0.28
Mato Grosso	0.31	0.21
Santa Catarina	0.27	0.19
Amazonas	0.09	0.03
Average	0.27	0.18
Groups		
Group I	0.30	0.21
Group II	0.40	0.20
Group III	0.31	0.22
Average	0.34	0.21

species and its genetic structure (Badfar-Chaleshtori et al. 2012), as well as their ability to estimate heterozygosity (Vogel et al. 1996). However, difficulties in relating fragment patterns for specific loci and genotypes in the genomes of polyploid species such as *D. trifida* and other tuberous plants limit the use of heterozygosity estimates to assess different molecular markers in these species (McGregor et al. 2000).

This study showed that it is possible to use both ISSR and SSR techniques for characterizing and discriminating morphologically distinct or similar yam accessions. Also, both ISSR and SSR results highlight the importance of traditional farmers in maintaining high genetic diversity among their local varieties. The eight SSR primer pairs were highly polymorphic and informative among the 53 *D. trifida* accessions analyzed in this study. The heterologous primers Da1A01 and Dab2C05 developed by Tostain et al. (2006) for *D. alata* L., *Dioscorea abyssinica* Hochst. ex Kunth and *Dioscorea praehensilis* Benth, showed 100 % polymorphism, and high discrimination power among accessions, equal to 0.97 and 0.91, respectively. These primers were also suitable for analyzing *D. trifida* accessions in the transferability tests conducted by Tostain et al. (2006). The primers specific for *D. trifida*, developed by Hochu et al. (2006), besides providing good resolution of bands in the gel electrophoresis, showed high polymorphism,

with an average of 93.8 %, and a high number of bands per primer (7 bands, on average). We highlight here primers MTI3 and MTI10 that revealed 13 and 11 bands, respectively (Table 2). In contrast, only five bands were revealed by primer MTI11, similar to results obtained by Hochu et al. (2006), which found only three bands for this primer.

Except for primers UBC 844, UBC 898 and DAT, which showed low polymorphism and low discrimination power among the accessions analyzed, all the other ISSR analyzed primers were highly polymorphic with a high number of bands per primer, especially for the UBC 7, UBC 898, JOHN and MANNY primers. These primers revealed more than 10 bands (Table 3), as well as high discrimination power among accessions, with  $D$  ranging from 0.36 to 0.82. The high percentage of polymorphism (75.8 %, on average) observed for ISSR was also reported by Zhou et al. (2008), analyzing 28 cultivars of *Dioscorea opposita* based on seven ISSR primers, which had a total of 65 fragments with 83 % polymorphism. High levels of polymorphism are common in ISSR, such as the results obtained for potatoes, with 90 % polymorphism (Prevost and Wilkinson 1999) and sweet potato, with 62.2 % (Huang and Sun 2000).

Mantel test results revealed that data obtained with the SSR and ISSR markers are correlated ( $r = 0.5$ ;  $p = 0.0002$ ), although they represent different genomic sequences. Several studies have also noted the existence of high correlation between different techniques of molecular markers in different species (Belaj et al. 2003; Biswas et al. 2010).

#### Genetic structure and conservation strategies

In Brazil, a center of diversity and domestication of various species, studies of genetic diversity are most often associated with economically important crops (Clement et al. 2010). Roots and tuber crops such as yams have been neglected by breeding and conservation research (Siqueira 2011). In this context, yam is considered an underutilized crop, subject to selection of interesting characters by the traditional communities, where farmers maintain varieties of their preference. Therefore, there is a low level of marketing and exchange of these materials when compared, for example, to other root crops, such as cassava, potato and sweet potato (Siqueira 2011). With this in mind, *D. trifida* can be affected by the process of genetic drift,



533 whose effect is intensified because of the presence of  
534 dioecy, which requires a male and a female plant in the  
535 same site for genetic recombination to occur between  
536 individuals (Mignouna et al. 2003). Although this  
537 species is allogamous in favorable climatic conditions,  
538 it is strongly maintained by vegetative propagation in  
539 cultivated fields, and in most cases, the clones of the  
540 same individuals are collected and planted for several  
541 years. However, the accessions analyzed in this study  
542 showed high genetic variability, with a variation in the  
543 similarity coefficient of Jaccard equal to 31 and 56 %  
544 for SSR and ISSR data, respectively. Veasey et al.  
545 (2012), by analyzing accessions of the same species  
546 collected in the Vale do Ribeira, São Paulo, based on  
547 isozymes, observed a variation in the similarity  
548 coefficient of Jaccard equal to 83 %.

549 In the cluster analysis, based on the Jaccard  
550 coefficient and UPGMA method, as well as in the  
551 PCoA and the Bayesian analyses, three genetically  
552 distinct and consistent groups were identified, with  
553 similar or identical membership. One of the three  
554 groups mentioned above consisted of accessions from  
555 Iguape municipality in the southern coast of the São  
556 Paulo State, which were grouped with accessions  
557 collected in the north coast of Santa Catarina, in the  
558 municipalities of Joinville and São Francisco do Sul. A  
559 second group was formed by accessions from the north  
560 coast of São Paulo, in Ubatuba municipality, and a  
561 third group classified the accessions collected in Mato  
562 Grosso, a region well apart from the others. The  
563 exception was the Amazonian accessions, purchased  
564 in local markets. Their group membership depended  
565 on the marker type or on the genomic region sampled.

566 The interesting fact in this study was the separation  
567 of accessions from the south and north coasts of São  
568 Paulo. Within Vale do Ribeira, in the south coast of  
569 São Paulo, Veasey et al. (2012) had already noticed a  
570 spatially structure in genetic variation along a much  
571 smaller geographic scale with isozyme markers.  
572 *D. trifida* local varieties from Vale do Ribeira were  
573 grouped according to their municipalities. Two clus-  
574 ters (with 100 % bootstrap) were obtained in the  
575 cluster analysis, one with varieties from Iguape  
576 municipality and the other with varieties from Cana-  
577 neia municipality. The same genetic structure was  
578 observed in the present study, but at a higher  
579 geographic scale, separating accessions among the  
580 north and south coastal areas in São Paulo State. The  
581 explanation for this finding is perhaps the introduction

of accessions of this species by waves of migrants  
from different regions or even different ethnic groups.  
On the other hand, there was a greater similarity of  
accessions on the south coast of São Paulo, Iguape,  
with those from Santa Catarina, suggesting that  
accessions from these two geographically adjacent  
regions have the same origin.

This dynamics can also be related to indigenous  
influence on the domestication of *D. trifida*, which has  
been cultivated by indigenous people from the coastal  
areas to the Central West region of Brazil (Pedralli  
1998). Among the indigenous groups involved in this  
process, we highlight the Guaraní, who are very  
itinerant and widely scattered throughout Brazil,  
including the coast of São Paulo (Ladeira 1992). This  
ethnic group traveled over long distances carrying  
with themselves various species of edible plants,  
among which *D. trifida* (Schmitz and Gazzaneo 1991).  
This species is, therefore, strongly associated with  
these indigenous people who, on the other hand, have a  
strong influence upon the traditional populations in the  
Vale do Ribeira, São Paulo (Veasey et al. 2012), where  
the Iguape municipality is located.

The maintenance of genetic variation is a major  
objective for conservation (Hamrick and Godt 1996)  
and knowledge of variation within and among popu-  
lations provides essential information in the formula-  
tion of appropriate conservation strategies (Francisco-  
Ortega et al. 2000). In this study, most of the genetic  
variability was observed within groups (60.6 and  
66.5 % for SSR and ISSR, respectively). In agreement  
with this result, in a study with wild and cultivated  
Guinea yams from south and south west Ethiopia,  
Mengesha et al. (2013) observed that most of the  
microsatellite diversity was found within rather than  
among populations. However, our study showed that  
even among groups the genetic variability was high  
(39.4 and 33.5 % for SSR and ISSR, respectively).  
Veasey et al. (2012), comparing two groups of  
*D. trifida* accessions analyzed with eight SSR primer  
pairs, one group including seven accessions from  
Iguape and Cananéia municipalities in São Paulo and  
one accession from Mato Grosso, and another group  
with four accessions from the Amazon, observed that  
most of the variation was between groups (62.9 %)  
compared with the variation within groups (37.1 %),  
in contrast to our findings. This result showed that the  
Amazonian accessions were genetically different from  
the other accessions.

In the present study, we observed a large genetic difference among accessions collected from the different locations analyzed, i.e., between groups, since the values of  $\Phi_{st}$  was 0.3937 for SSR and 0.3353 for ISSR, which corresponds to a low gene flow among the regions studied. This pattern demonstrates that, unlike other species grown in traditional agriculture, such as cassava (Siqueira et al. 2009) or sweet potato (Veasey et al. 2008), *D. trifida* is a more regionalized crop or less scattered when compared to other yam species, such as *D. alata* (Bressan et al. 2011; Siqueira et al. 2012). This observation can be related to historical and socioeconomic factors, such as the different ways of using these tubers and variation in the preference of varieties over time and space (Veasey et al. 2010). The positive correlation between genetic and geographical distances obtained in the Mantel test, confirms the structure of these materials in the geographic area sampled. Although the value of this correlation was low, the fact that it is significant at 1 % confirms the spatially structure of the genetic variation, which is consistent with the observed  $\Phi_{st}$  value. However, the low correlation values could be due to possible recent exchange of materials among these regions, carried out by non-governmental organizations working with indigenous groups, to encourage the exchange of genotypes and inserting new material to prevent genetic erosion of these varieties. Thus, exchange fairs are held, allowing the acquisition and supply of new genotypes. There is little information about these indigenous groups, such as the Guaraní group mentioned above, who have survived various pressures, such as land struggles, beyond sociocultural pressures of modern societies (Arruda 1999). These difficulties are common to both indigenous groups and for farmers that grow various species in a traditional way in the tropics.

Considering the genetic differences among the genotypes grown in the studied locations, the accessions of São Paulo showed higher diversity, followed by Mato Grosso, Santa Catarina and Amazonas. The average level of genetic diversity present in accessions of *D. trifida* was 0.27 and 0.18 among states, for SSR and ISSR data, respectively. Considering the groups formed in the Bayesian analysis, using Structure software, higher diversity was found among accessions in group II for SSRs, which includes accessions from three states (São Paulo, Santa Catarina and Amazonas). However, for the ISSR marker, the

highest diversity was found for group III (among Mato Grosso accessions), although the values were very similar among the three groups. In general, the average level of genetic diversity was 0.34 and 0.21, for SSR and ISSR data, respectively. These indices are greater than the values found for perennial herbs ( $H' = 0.17$ ) and to those species with wide geographic distribution ( $H' = 0.20$ ) (Hamrick and Godt 1989). However, studies by Zhou et al. (2008) with varieties of *D. opposita*, widely used in Chinese medicine, showed  $H' = 0.32$ , a relatively high value and in most cases superior to that obtained for *D. trifida* in this study. It is noteworthy that several factors influence the level of genetic diversity of a species, among them, the geographic distribution, life cycle, reproductive system, dispersal patterns, population size, among others (Hamrick et al. 1991; Gaudeul et al. 2000; Zhou et al. 2008).

The level of genetic diversity observed among accessions is directly related to the fact that *D. trifida* is a polyploid species and reproduces both by outcrossing and vegetative propagation. Thus, individuals are usually highly heterozygous, preserving the allelic diversity at the individual level (Veasey et al. 2008; Siqueira et al. 2009). However, the emergence of variant plants arising from seeds, result of genetic recombination, is unlikely, since the occurrence of flowering and fruiting in this species was not reported and detected by farmers during plant collection.

The genetic diversity level displayed by the accessions collected in the States of São Paulo, Mato Grosso and Santa Catarina was expected because of the wide distribution of the crop in these regions. However, we also found genetically similar varieties, which can be related to the fact that the collection has been restricted to a few farmers and a few communities in Ubatuba, SP, where practically all the collection was made in the community of the Sertão de Ubatumirim, and in the community Pirabeiraba in Santa Catarina. The similarity between accessions corroborates the names given by farmers in the studied regions. In São Paulo, except for accession 195, called *cara cobrinha* (little snake yam), varieties were obtained with only two popular names, *cara roxo* (purple yam) and *cara branco* (white yam), the latter being represented in smaller numbers. In Mato Grosso, most accessions also received the name *cara roxo*. In Santa Catarina, the predominant designations for *D. trifida* were *cara* and *cara mimoso*.

In conclusion, our results suggest that both markers were useful for evaluation of genetic diversity and assessing differentiation between *D. trifida* populations in Brazil. However, the SSR marker detected higher diversity indices while the ISSR marker seemed more efficient in the clustering of the different genotypes, being able to separate the two Amazonian accessions in the cluster analysis. But both markers detected high levels of genetic diversity for accessions of *D. trifida* maintained by traditional farmers in the states of São Paulo, Mato Grosso, Santa Catarina and Amazonas. The high within-group variation found is quite interesting for the maintenance of the crop over time. Knowledge of the genetic relationships among accessions is an important information for the efficient use and conservation of this species, both *ex situ* in genebanks, or *in situ*, within the aim of conservation in the rural property, known as on-farm conservation. Considering that high genetic diversity was found both within and between groups of accessions from different regions visited, the collection and conservation strategies should consider a large number of individuals from all regions sampled in order to cover all the genetic diversity present in these materials. On-farm conservation, in the case of *D. trifida*, is quite interesting for considering the socio-cultural factors involved in the evolution of the species, considering that agricultural practices, through the cultivation and artificial selection, allows the accumulation over time of morphological traits of agronomic interest, providing an enrichment of genetic variability not only of this species, but of other crops as well.

**Acknowledgments** The authors would like to thank the researchers Nivaldo Peroni, Antônio Henrique dos Santos, Lin Chau Ming, Edson Ferreira da Silva, Marcos V.B.M. Siqueira and Almecina Balbino Ferreira for their assistance in this research and the agriculturists for their contributions in the field collecting and interviews. The authors would also wish to thank FAPESP (process no. 2007/04805-2) and CNPq for the financial support given to this study.

## References

- Arruda R (1999) "Populações tradicionais" e a proteção dos recursos naturais em unidades de conservação. *Ambient Soc* 5:79–252
- Ayensu ES, Coursey DG (1972) Guinea yams: the botany, ethnobotany, use and possible future of yams in West Africa. *Econ Bot* 26:301–318

- Badfar-Chaleshtori S, Shiran B, Kohgard M, Mommeni H, Hafizi A, Khodambashi M, Mirakhorli N, Sorkheh K (2012) Assessment of genetic diversity and structure of Imperial Crown (*Fritillaria imperialis* L.) populations in the Zagros region of Iran using AFLP, ISSR and RAPD markers and implications for its conservation. *Biochem Syst Ecol* 42:35–48
- Belaj A, Satovic Z, Cipriani G, Baldoni L, Testolin R, Rallo L, Trujillo I (2003) Comparative study of the discriminating capacity of RAPD, AFLP and SSR markers and of their effectiveness in establishing genetic relationships in olive. *Theor Appl Genet* 107:736–744
- Biswas MK, Xu Q, Deng X (2010) Utility of RAPD, ISSR, IRAP and REMAP markers for the genetic analysis of *Citrus* spp. *Sci Hortic* 124:254–261
- Bousalem M, Amau G, Hochu I, Amolin R, Viader V, Santoni S, David J (2006) Microsatellite segregation analysis and cytogenetic evidence for tetrasomic inheritance in the American yam *Dioscorea trifida* and a new basic chromosome number in the *Dioscoreae*. *Theor Appl Genet* 113:439–451
- Bousalem M, Viader V, Mariac C, Gomez R, Hochu I, Santoni S, David J (2010) Evidence of diploidy in the wild Amerindian yam, a putative progenitor of the endangered species *Dioscorea trifida* (Dioscoreaceae). *Genome* 53:371–383
- Bressan EA, Veasey EA, Peroni N, Felipim AP, Santos KMP (2005) Collecting yam (*Dioscorea* spp.) and sweet potato (*Ipomoea batatas*) germplasm in traditional agriculture small-holdings in the Vale do Ribeira, São Paulo, Brazil. *Plant Genet Resour Newsl* 144:8–13
- Bressan EA, Briner Neto T, Zucchi MI, Rabello RJ, Veasey EA (2011) Morphological variation and isozyme diversity in *Dioscorea alata* L. landraces from Vale do Ribeira, Brazil. *Sci Agric* 68:494–502
- Clement CR (1999) 1492 and the loss of Amazonian crop genetic resources. II. Crop biogeography at contact. *Econ Bot* 53:203–216
- Clement CR, Cristo-Araújo M, D'Eeckenbrugge GC, Pereira AA, Picanço-Rodrigues D (2010) Origin and domestication of native Amazonian crops. *Diversity* 2:72–106
- Coursey DG (1976) The origin and domestication of yams in Africa. In: Harlan JR, De Wet JMJ, Stemler ABL (eds) *Origin of African plant domestication*. Mouton Hague, Netherlands, pp 383–408
- Creste S, Tulmann NA, Figueira A (2001) Detection of single sequence repeat polymorphisms in denaturing polyacrylamide sequencing gels by silver staining. *Plant Mol Biol Rep* 19:299–306
- Degras L (1993) *The yam: a tropical root crop*. Macmillan Press, London
- Dickau R, Ranere AJ, Cooke RG (2007) Starch grain evidence for the preceramic dispersals of maize and root crops into tropical dry and humid forests of Panama. *Proc Natl Acad Sci* 104:3651–3656
- Doyle JJ, Doyle JL (1990) Isolation of plant DNA from fresh tissue. *Focus* 12:13–15
- Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Mol Ecol* 14:2611–2620
- Falush D, Stephens M, Pritchard JK (2007) Inference of population structure using multilocus genotype data: dominant markers and null alleles. *Mol Ecol Notes* 7:574–578

- 837 Francisco-Ortega J, Santos-Guerra A, Kim SC, Crawford DJ  
838 (2000) Plant genetic diversity in the Canary Islands: a  
839 conservation perspective. *Am J Bot* 87:909–919
- 840 Gaudeul M, Taberlet P, Till-bottraud I (2000) Genetic diversity  
841 in an endangered alpine plant, *Eryngium alpinum* L.  
842 (Apiaceae), inferred from amplified fragment length  
843 polymorphism markers. *Mol Ecol* 9:1625–1637
- 844 Hamrick JL, Godt MJW (1989) Allozyme diversity in plant  
845 species. In: Brown AHD, Clegg MT, Kahler AL, Weir BS  
846 (eds) Plant population genetics, breeding and genetic  
847 resources. Sinauer Associates, Massachusetts, pp 43–46
- 848 Hamrick JL, Godt MJW (1996) Conservation genetics of  
849 endemic plant species. In: Avise JC, Hamrick JL (eds)  
850 Conservation genetics: case histories from nature. Chap-  
851 man and Hall, New York, pp 281–304
- 852 Hamrick JL, Godt MJ, Murawski DA, Loveless MD (1991)  
853 Correlations between species and allozyme diversity:  
854 implications for conservation biology. In: Falk DA, Hol-  
855 singer KE (eds) Genetics and conservation of rare plants.  
856 Oxford University Press, New York, pp 75–86
- 857 Hochu I, Santoni S, Bousalem M (2006) Isolation, character-  
858 ization and cross-species amplification of microsatellite  
859 DNA loci in the tropical American yam *Dioscorea trifida*.  
860 *Mol Ecol* 6:137–140
- 861 Huang JC, Sun M (2000) Fluorescein PAGE analysis of  
862 microsatellite-primed PCR: a fast and efficient approach  
863 for genomic fingerprinting. *Biotechniques* 28:1069–1072
- 864 Ladeira MI (1992) “O caminhar sob a luz”-o território Mbyá à  
865 beira do Oceano. Pontifícia Universidade Católica de São  
866 Paulo, Dissertation
- 867 Lebot V (2009) Tropical root and tuber crops: cassava, sweet  
868 potato, yams and aroids. CABI, London
- 869 Mantel N (1967) The detection of disease clustering and a  
870 generalized regression approach. *Cancer Res* 27:202–209
- 871 Mattioni C, Casasoli M, Gonzalez M (2002) Comparison of  
872 ISSR and RAPD markers to characterize three Chilean  
873 Nothofagus species. *Theor Appl Genet* 104:1064–1070
- 874 McGregor CE, Lambert CA, Greyling MM, Louw JH, Warnich  
875 L (2000) A comparative assessment of DNA fingerprinting  
876 techniques (RAPD, ISSR, AFLP and SSR) in tetraploid  
877 potato (*Solanum tuberosum* L.) germplasm. *Euphytica*  
878 113:135–144
- 879 Mengesha WA, Demissew S, Fay MF, Smith RJ, Nordal I,  
880 Wilkin P (2013) Genetic diversity and population structure  
881 of Guinea yams and their wild relatives in South and South  
882 West Ethiopia as revealed by microsatellite markers. *Genet*  
883 *Resour Crop Evol* 60:529–541
- 884 Mignouna HD, Abang MM, Fagbemi SA (2003) A comparative  
885 assessment of molecular marker assays (AFLP, RAPD and  
886 SSR) for white yam (*Dioscorea rotundata*) germplasm  
887 characterization. *Ann Appl Biol* 142:269–276
- 888 Milbourne D, Meyer R, Collins AJ, Ramsay LD, Gebhardt C,  
889 Waugh R (1998) Isolation, characterisation and mapping of  
890 simple sequence repeat loci in potato. *Mol Gen Genet*  
891 259:233–245
- 892 Montaldo A (1991) Cultivo de raíces y tubérculos tropicales.  
893 Instituto Interamericano de Ciencias Agrícolas de la OEA,  
894 Lima
- 895 Odu BO, Asiedu R, JA Hughes, Shoyinka SA, Oladiran OA  
896 (2004) Identification of resistance to yam mosaic virus  
(YMV), genus *Potyvirus* in white Guinea yam (*Dioscorea*  
*rotundata* Poir). *Field Crop Res* 89:97–195
- Oliveira EJ, Pádua JG, Zucchi MI, Vencovsky R, Vieira LC  
(2006) Origin, evolution and genome distribution of  
microsatellites. *Genet Mol Biol* 29:294–307
- Olsen KM (2004) SNPs, SSRs and inferences on cassava’s  
origin. *Plant Mol Biol* 56:517–526
- Pedralli G (1998) Revisão taxonômica das espécies de Dio-  
scoreaceae (R.Br.) Lindley da Cadeia do Espinhaço, Minas  
Gerais e Bahia, Brasil. Dissertation, University of São  
Paulo
- Perrier X, Jacquemoud-Collet JP (2006) DARwin software.  
<http://darwin.cirad.fr/darwin>. Accessed 20 Nov 2011
- Piperno DR, Ranere AJ, Holst I, Hansell P (2000) Starch grains  
reveal early root crop horticulture in the Panamanian  
tropical forest. *Nature* 407:894–897
- Prevost A, Wilkinson MJ (1999) A new system of comparing  
PCR primers applied to ISSR fingerprinting of potato  
cultivars. *Theor Appl Genet* 98:107–112
- Pritchard JK, Donnelly P (2001) Case-control studies of asso-  
ciation in structured or admixed populations. *Theor Popul*  
*Biol* 60:227–237
- Pritchard JK, Stephens M, Donnelly P (2000) Inference of  
population structure using multilocus genotype data.  
*Genetics* 155:945–959
- Ramos-Escudero F, Santos-Buelga C, Pérez-Alonso JJ, Yáñez  
JA, Dueñas M (2010) HPLC-DAD-ESI/MS identification  
of anthocyanins in *Dioscorea trifida* L. yam tuber (purple  
sachapapa). *Eur Food Res Technol* 230:745–752
- Reddy MP, Sarla N, Reddy EA (2002) Inter simple sequence  
repeat (ISSR) polymorphism and application plant breed-  
ing. *Euphytica* 128:9–17
- Rogstad SH (1992) Saturated NaCl-CTAB solution as a means  
of field preservation of leaves for DNA analyses. *Taxon*  
41:701–708
- Rohlf FJ (1992) NTSYS-pc: numerical taxonomy and multi-  
variate analysis system, version 1.70 (software). Stony  
Brook, New York
- Schmitz PI, Gazzaneo M (1991) O que comia o Guarani pré-  
colonial. *Rev Arqueol* 6:89–105
- Schneider S, Roessli D, Excoffier L (2000) Arlequin: software  
for population data analysis (Software). Genetic and  
Biometry laboratory of University of Geneva, Geneva
- Siqueira MVBM (2011) Yam: a neglected and underutilized  
crop in Brazil. *Hort Brasil* 29:6–20
- Siqueira MVBM, Queiroz-Silva JR, Bressan EA, Borges A,  
Pereira KJC, Pinto JG, Veasey EA (2009) Genetic char-  
acterization of cassava (*Manihot esculenta*) landraces in  
Brazil assessed with simple sequence repeats. *Genet Mol*  
*Biol* 32:104–110
- Siqueira MVBM, Dequigiovanni G, Corazon-Guivin M, Feltran  
J, Veasey EA (2012) DNA fingerprinting of water yam  
(*Dioscorea alata*) cultivars in Brazil based on microsatel-  
lite markers. *Hort Brasil* 30:653–659
- Stephens JM (2009) Cushcush – *Dioscorea trifida* L.  
<http://edis.ifas.ufl.edu/my057>. Accessed 27 Dec 2009
- Tessier C, David J, This P, Boursiquot J, Charrier A (1999)  
Optimizations of the choice of molecular markers for  
varietal identification in *Vitis vinifera* L. *Theor Appl Genet*  
98:171–177

- 957 Tostain S, Scarcelli N, Brottier P, Marchand JL, Pham JL, Noyer  
958 JL (2006) Development of DNA microsatellite markers in  
959 tropical yam (*Dioscorea* sp.). *Mol Ecol* 6:173–175
- 960 Veasey EA, Borges A, Rosa MS, Queiroz-Silva JR, Bressan EA,  
961 Peroni N (2008) Genetic diversity in Brazilian sweetpotato  
962 (*Ipomoea batatas* (L.) Lam) landraces assessed with  
963 microsatellites. *Genet Mol Biol* 31:725–733
- 964 Veasey EA, Monteiro MVB, Gomes LR, Nascimento WF,  
965 Ferreira AB, Silva MS, Silva EF, Ming LC, Peroni N,  
966 Santos AH (2010) Ocorrência e diversidade de espécies  
967 cultivadas do gênero *Dioscorea* em diversos agroecos-  
968 sistemas brasileiros. In: Ming LC, Amorozo MCM, Kffuri  
969 CW (eds) *Agrobiodiversidade no Brasil: experiências e*  
970 *caminhos da pesquisa*, 1st edn. NUPEEA, Recife, pp 45–74
- 971 Veasey EA, Bressan EA, Siqueira MVB, Borges A, Queiroz-  
972 Silva JR, Pereira KJC, Recchia GH, Ming LC (2012)  
973 Genetic characterization of cassava (*Manihot esculenta*  
974 Crantz) and yam (*Dioscorea trifida* L.) landraces in swi-  
975 den agriculture systems in Brazil. In: Gepts P, Famula TR,  
976 Bettinger RL, Brush SB, Damania AB, McGuire PE,  
977 Qualset CO (eds) *Biodiversity in agriculture: domestica-*  
978 *tion, evolution, and sustainability*. Cambridge Univ. Press,  
979 New York, pp 344–360
- 980 Velez GA (1998) The chagra: collective patrimony of the  
981 indigenous Amazonian communities. *Beyond Law* 6:  
982 121–142
- Vogel JM, Rafalski A, Powell W, Morgante M, Andre C, 983  
Hanafey M, Tingey SV (1996) Application of genetic 984  
diagnostics to plant genome analysis and plant breeding. 985  
*HortScience* 31:1107–1108 986
- Wolfe AD (2000) ISSR protocols. [http://www.biosci.ohio-sta-  
te.edu/~awolfe/ISSR/protocols.ISSR.html](http://www.biosci.ohio-state.edu/~awolfe/ISSR/protocols.ISSR.html). Accessed 18 987  
May 2009 988
- Wolfe AD (2005) ISSR techniques for evolutionary biology. 990  
*Methods Enzymol* 395:134–144 991
- Wu Z, Leng C, Tao Z, Wei Y, Jiang C (2009) Genetic diversity 992  
of *Dioscorea alata* on ISSR analysis. *Zhongguo Zhong* 993  
*Yao Za Zhi* 34:3017–3020 994
- Yeh FC, Yang RC, Boyle TBJ, Ye ZH, Mao JX (1997) POP- 995  
GENE, the user-friendly shareware for population genetic 996  
analysis. Mol Biol Biotechnol Centre, Canadian, Edmonton 997
- Zhou Y, Zhou C, Yao H, Liu Y, Tu R (2008) Amplification of 998  
ISSR markers in detection of genetic variation among 999  
Chinese yam (*Dioscorea opposita* Thunb) cultivars. *Life* 1000  
*Sci J* 5:6–12 1001
- Zietkiewicz E, Rafalski A, Labuda D (1994) Genome finger- 1002  
printing by simple sequence repeat (SSR)-anchored poly- 1003  
merase chain reaction amplification. *Genomics* 20:176–183 1004  
1005

UNCORRECTED