

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

Photoelectron Spectroscopy of Rare Gas Dimers Revisited: Vibrationally Resolved Photoelectron Spectrum of Argon Dimer

### Permalink

<https://escholarship.org/uc/item/9hv0n9tf>

### Journal

Journal of Chemical Physics, 98(7)

### Authors

Pradeep, T.

Niu, B.

Shirley, D.A.

### Publication Date

1992-11-01



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

## CHEMICAL SCIENCES DIVISION

Submitted to Journal of Chemical Physics

### Photoelectron Spectroscopy of Rare Gas Dimers Revisited: Vibrationally Resolved Photoelectron Spectrum of Argon Dimer

T. Pradeep, B. Niu, and D.A. Shirley

November 1992



LOAN COPY |  
Circulates |  
for 4 weeks | Bldg. 50 Library.  
Copy 2

LBL-32925

### DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. Neither the United States Government nor any agency thereof, nor The Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or The Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or The Regents of the University of California and shall not be used for advertising or product endorsement purposes.

Lawrence Berkeley Laboratory is an equal opportunity employer.

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

LBL -32925  
UC-401

**Photoelectron spectroscopy of rare gas dimers  
revisited: vibrationally resolved photoelectron  
spectrum of argon dimer**

T. Pradeep<sup>1</sup>, B. Niu and D. A. Shirley<sup>2</sup>

Chemical Sciences Division,  
Lawrence Berkeley Laboratory,  
University of California,  
One Cyclotron Road, Berkeley,  
CA 94720, USA.

November 1992

This work was supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Chemical Sciences Division, of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098.

**ABSTRACT**

Photoelectron spectra of rare gas dimers  $\text{Ar}_2$ ,  $\text{Kr}_2$  and  $\text{Xe}_2$  have been measured using the HeI radiation at 584 Å with a resolution of 13 meV. All the six ionic states in the HeI region have been resolved for the first time, including some of the purely repulsive states. Ionization energies are presented to an accuracy  $\leq 0.003$  eV. Dissociation energies ( $D_0$ ) of the ionic states are calculated using the best estimate adiabatic ionization energies. The  $D^2\Sigma^+_{1/2g}$  states of all the dimers are weakly bound, contrary to an earlier theoretical prediction. Part of the vibrational structure of two of the electronic states of  $\text{Ar}_2$  has been resolved. For the  $A^2\Sigma^+_{1/2u}$  state as many as 14 vibrational excitations are observed. These excitations are assigned to transitions to the higher vibrational levels of  $v = 32 - 45$ . For the  $B^2\Pi_{3/2g}$  state, three vibrational structures are resolved and are assigned to  $v = 2 - 4$  of the ionic state. Accurate values of the spectroscopic constants of these states are presented. The present values are compared with the values available in the literature. The  $D_e$  values for the  $A^2\Sigma^+_{1/2u}$  and the  $B^2\Pi_{3/2g}$  states are estimated to be 1.361 and 0.104 eV respectively.

## I. INTRODUCTION

Dissociation energies of rare gas dimers are much smaller than ambient thermal energy and consequently equilibrium methods are not adequate to study them. Since the development of molecular beam methods, there have been numerous attempts to study a variety of properties of these fascinating species. Part of the interest of late was due to the possible use of these molecules in vacuum ultraviolet lasers. In fact, rare gas dimers are the most studied class of van der Waals molecules. Electronic structure of these species have been studied both by theory<sup>1-3</sup> and by experiment<sup>4-10</sup>. The neutral dimers have a weak van der Waals minimum at a large internuclear distance. However, the ionic states are strongly bound. For example, while the ground state of  $\text{Ar}_2^+$  has a dissociation energy<sup>4</sup> ( $D_e$ ) of over  $10000 \text{ cm}^{-1}$ , the neutral dimer<sup>11</sup> is bound by only  $98.7 \text{ cm}^{-1}$ .

There are six electronic states for the rare gas dimer ions in the HeI energy region. Adiabatic IEs of all the dimers are much lower than the ionization potentials of the atoms, implying that the ground state dimer ions are strongly bound. HeI photoelectron spectroscopic studies by Dehmer and Dehmer<sup>4,5</sup> showed that many of the excited ionic states are also attractive. However, they could not observe all the ionic states. Although some of the states are strongly bound, no vibrational structure was observed in the photoelectron spectra. Resonantly enhanced multiphoton ionization was also carried out to study the photoelectron spectrum<sup>10</sup> of  $\text{Xe}_2$ . In addition to the previously observed states, the  $\text{C}^2\Pi_{3/2u}$  and the  $\text{D}^2\Sigma^+_{1/2g}$  states were observed and found to be weakly bound. This was the only photoelectron spectroscopic study in which all the ionic states were observed.

The present study of the photoelectron spectra of rare gas dimers was undertaken with the objectives of: 1) obtaining the ionization energies of all the

electronic states, particularly the  $C^2\Pi_{3/2u}$  and the  $B^2\Pi_{1/2g}$  states, which have eluded most of the previous attempts and 2) resolving the vibrational structure of at least a few more strongly bound states. We have been successful in both these objectives.

The vibrational structure of the ionic states has been observed before. Coincidence between threshold electrons and ions was used<sup>12</sup> to determine the vibrational structure of the ground state of  $Ar_2^+$ . That study gave a  $\omega_e$  value of  $310.8\text{ cm}^{-1}$  for this state. A progression beginning at  $v = 3$  was observed and as many as 20 vibrational excitations were seen in the spectrum. Vibrational spectroscopy of  $Xe_2^+$  was also carried out by pulsed field ionization<sup>13</sup>. For the  $A^2\Sigma^+_{1/2u}$  state, a progression beginning at  $v = 56$  was observed. Although vibrational excitations were observed for the  $B^2\Pi_{3/2g}$  state, absolute assignments were not made. For the  $C^2\Pi_{3/2u}$  state, vibrational excitations to the  $v = 0 - 2$  levels were observed. To the best of our knowledge, vibrational spectroscopy has not been carried out on the ionic states of  $Kr_2$ .

In this paper, we report the photoelectron spectroscopic study of the rare gas dimers  $Ar_2$ ,  $Kr_2$  and  $Xe_2$ . All the six ionic states in the HeI region have been observed, including some of the purely repulsive states. Although our values of the adiabatic IEs are only best estimates, accurate values of the vertical IEs are presented. Vibrational structure has been resolved in two electronic states of  $Ar_2$ , giving precise values of  $\omega_e$  and  $D_e$ . Experimental parameters are compared with existing values in the literature.

## II. EXPERIMENT

The molecular beam photoelectron spectrometer used in the present investigation has been fully described elsewhere<sup>14</sup> and only a brief description is provided here. Supersonic beams of the unseeded rare gases from a  $70\ \mu$  nozzle skimmed by a 0.9 mm skimmer were crossed by the HeI radiation at  $90^\circ$ . The photoelectrons were sampled at  $90^\circ$  with respect to both the atomic and the photon



beams and were analyzed by a hemispherical sector at an operating pass energy of 1 eV. The operating resolution was 13 meV as measured on Ar  $2P_{3/2}$ . The spectra were measured at different stagnation pressures and nozzle temperatures. The exact conditions vary for each rare gas dimer studied. The conditions were optimized to obtain the spectrum essentially devoid of the heavier clusters. The resolution degraded to 14 meV during the day-long experiments.

### III. RESULTS AND DISCUSSION

In the linear combination of atomic orbitals (LCAO) description, the single determinant configuration of the rare gas dimer is,

$$\dots(\sigma_g ns)^2 (\sigma_u ns)^2 (\sigma_g np)^2 (\pi_u np)^4 (\pi_g np)^4 (\sigma_u np)^2$$

giving a  $1\Sigma_g^+$  ground state. The value of  $n$  is 3, 4 and 5 for Ar<sub>2</sub>, Kr<sub>2</sub> and Xe<sub>2</sub> respectively. Since the bond order is zero, the dimers have only polarization minima. The well depth is 12, 17 and 24 meV for Ar<sub>2</sub>, Kr<sub>2</sub> and Xe<sub>2</sub> respectively<sup>11,15</sup>. The ionic states to be observed in the HeI region are,

$$\dots(\sigma_g ns)^2 (\sigma_u ns)^2 (\sigma_g np)^2 (\pi_u np)^4 (\pi_g np)^4 (\sigma_u np)^1 \quad A^2\Sigma_u^+$$

$$\dots(\sigma_g ns)^2 (\sigma_u ns)^2 (\sigma_g np)^2 (\pi_u np)^4 (\pi_g np)^3 (\sigma_u np)^2 \quad B^2\Pi_g$$

$$\dots(\sigma_g ns)^2 (\sigma_u ns)^2 (\sigma_g np)^2 (\pi_u np)^3 (\pi_g np)^4 (\sigma_u np)^2 \quad C^2\Pi_u^+$$

$$\dots(\sigma_g ns)^2 (\sigma_u ns)^2 (\sigma_g np)^1 (\pi_u np)^4 (\pi_g np)^4 (\sigma_u np)^2 \quad D^2\Sigma_g^+$$

in the order of increasing ionization energy.

The A and B states arise out of the removal of antibonding electrons and thus are attractive. The C and D states are the result of removal of bonding electrons and thus produce repulsive wells. As far as the relative stability of A and B states is concerned, the A state is more attractive than the B state since  $\sigma_u np$  is more antibonding than  $\pi_g np$ . Spin-orbit effects will split these states into six states. In this nomenclature, the states to be observed in the HeI region are  $A^2\Sigma^+_{1/2u}$ ,  $B^2\Pi_{3/2g}$ ,

$C^2\Pi_{3/2u}$ ,  $B^2\Pi_{1/2g}$ ,  $C^2\Pi_{1/2u}$  and  $D^2\Sigma^+_{1/2g}$ , in order of increasing ionization energy. The first four states dissociate into  $(Rg) ^1S_0$  and  $(Rg^+) ^2P_{3/2}$  and the remaining two to  $(Rg) ^1S_0$  and  $(Rg^+) ^2P_{1/2}$ . These states are designated 1, 2, 3, 4, 5, and 6 in the text. Although the calculations of Wadt<sup>2</sup> show a curve crossing between the  $1/2g$  and  $3/2u$  states at an internuclear distance of 3.25 - 3.50 Å, this is not important in the present experiment since this distance is considerably shorter than the equilibrium internuclear distance and is far away from the Franck-Condon region.

Adiabatic IEs of the rare gas dimers have been determined by various workers. The best values of the IEs are  $14.44 \pm 0.02$  ( $Ar_2$ )<sup>6</sup>,  $12.87 \pm 0.015$  ( $Kr_2$ )<sup>7</sup> and  $11.13 \pm 0.01$  ( $Xe_2$ )<sup>8</sup> eV. Photoionization studies of Ar clusters<sup>16</sup> give a different value for the adiabatic IE of  $Ar_2$ , which is 0.05 eV higher than the above estimate. Because of their small Franck-Condon overlap, these adiabatic IEs have not been observed in the photoelectron spectra. For example, the appearance energy observed in photoelectron spectroscopy<sup>4</sup> for  $Ar_2$  is 15.22 eV, higher by 0.78 eV than the adiabatic IE.

The dimers are discussed separately below.

#### A. $Ar_2$

Photoelectron spectra of all the rare gas dimers are essentially similar. In Fig. 1 we show the photoelectron spectrum of a mixture of Ar and  $Ar_2$  obtained at a stagnation pressure of 280 torr and a nozzle temperature of 80 K. The dimer ion features are indicated by numbers 1, 2, 4, 5 and 6 in the figure. Features 1 and 2 are well separated from the intense atomic line, which is 100 times intense than the dimer features. Peak 5 is also well separated from the atomic line. All these features are shown up in the photoelectron spectrum of Dehmer and Dehmer<sup>4</sup> also. These dimer features show strong stagnation pressure dependence. As reported by Dehmer and Dehmer<sup>4</sup>, a strong pressure dependence was exhibited by peak 5. Below a stagnation

pressure of 100 torr no dimer features were visible at a nozzle temperature of 300 K. The features due to the dimer increased in intensity up to a stagnation pressure of 800 torr. Above this pressure, higher clusters were also observed. A gradual increase in the intensity below the appearance energy of the dimer is attributed to the presence of heavier clusters. The evolution of the spectra was studied both by increasing the stagnation pressure and by decreasing the nozzle temperature. As the pressure increased above 800 torr, the appearance energy kept decreasing. However, the dimer features, particularly the peak 5, was still visible. The evolution of the spectra did not suggest the presence of ionization chromophores unlike that reported by Carnovale et al<sup>17</sup> who suggest  $\text{Ar}_{13}$  to be the ionization chromophore of heavier clusters and condensed Ar. According to Hegena's scaling laws<sup>18</sup>, in the case of Ar for a nozzle diameter of 70  $\mu$ , a nozzle temperature of 80 K, and a stagnation pressure of 500 torr, the scaling parameter  $\Gamma^*$  is about 1700, which corresponds to an average cluster size<sup>19</sup> of 100. The spectrum at these conditions did not show features similar to  $\text{Ar}_{13}$ . The spectrum showed only a broad unstructured band extending over an ionization energy of 1.5 eV below the atomic  $^2\text{P}_{3/2}$  peak. Moreover, the spectra of other rare gas clusters  $\text{Kr}_n$  and  $\text{Xe}_n$  also did not exhibit features suggesting the existence of ionization chromophores.

The spectrum at the lowest stagnation pressure showed only features due to the impurity lines in the undispersed HeI light source, apart from the intense atomic peaks. The features labeled 1, 2, 4, 5 and 6 in Fig. 1 are attributed to the dimer, by comparison with the spectrum of Dehmer and Dehmer<sup>4</sup> and with other rare gas dimer spectra, and on the basis of the stagnation pressure dependence of the features. Feature 1 is assigned to the ground ionic state. The appearance energy of the ion is  $15.138 \pm 0.003$  eV, below which the intensity is at the background level at  $\sim 1$  c/s. This appearance energy is lower than the value of 15.22 eV determined by Dehmer and

Dehmer<sup>4</sup> and higher by 0.698 eV than the adiabatic IE. This difference is due to the poor Franck-Condon (FC) overlap between the two states as mentioned earlier. This feature shows distinct vibrational features. As many as 9 features are clearly visible in Fig. 1. The features have been observed repeatedly in separate scans and also at different stagnation pressures, always appearing at the same ionization energies with the same intensity pattern. The vertical IE value of this state is  $15.548 \pm 0.003$  eV.

Feature 2 is assigned to the  $B^2\Pi_{3/2g}$  state. This state, with a dissociation energy of  $\sim 0.1$  eV, is expected to show vibrational structure at high resolution. A careful examination of the feature shows three distinct peaks at ionization energies 15.673(1), 15.689(1) and 15.702(1) eV. The extended vibrational structure of the  $A^2\Sigma^+_{1/2u}$  ground ionic state overlaps with this state, and one or more vibrational excitations of the  $B^2\Pi_{3/2g}$  state are unresolved. The adiabatic and vertical ionization energies are  $15.630 \pm 0.003$  and  $15.689 \pm 0.001$  eV respectively. Our values are in excellent agreement with those of Dehmer and Dehmer<sup>4</sup>.

The second excited state of the ion ( $C^2\Pi_{3/2u}$ ) has not been observed in previous studies. This state was thought to be obscured by the intense atomic  $^2P_{3/2}$  line. However, a peak is visible in Fig. 2, where a semi-log plot of the energy region of interest is presented. A shoulder in the low ionization energy side of the atomic  $^2P_{3/2}$  peak at 15.734(3) eV is assigned to the  $C^2\Pi_{3/2u}$  state on the basis of the considerations discussed above. Observation of this feature at the low IE side of the atomic  $^2P_{3/2}$  peak indicates that this state is weakly bound as predicted by Wadt<sup>2</sup>. Our best estimate of the adiabatic IE of this state is  $15.708 \pm 0.003$  eV. This is the first experimental observation of the  $C^2\Pi_{3/2u}$  state.

Peak 4 is assigned to the  $B^2\Pi_{1/2g}$  state. Although this feature is visible in the spectrum of Dehmer and Dehmer<sup>4</sup> they left it unassigned. The stagnation pressure dependence of this feature makes us believe that it is indeed due to the dimer.

Observation of this feature at the higher IE side of the atomic line suggests that the  $B^2\Pi_{3/2g}$  state is repulsive in the FC region in agreement with Wadt's prediction<sup>2</sup>.

Peak 5 is assigned to the  $C^2\Pi_{1/2u}$  state. This is the most intense feature due to the dimer in the spectrum and shows pronounced stagnation pressure dependence. It is one of the characteristic peaks due to the dimer persisting even at very low stagnation pressures. The values of its adiabatic and vertical IEs are  $15.847 \pm 0.003$  and  $15.877 \pm 0.001$  eV respectively.

Peak 6 corresponding to the  $D^2\Sigma^+_{1/2g}$  state, is partly obscured by the more intense Ar  $^2P_{3/2}$  (HeI 591.41 Å) peak. Our adiabatic and vertical IE estimates are  $15.934 \pm 0.003$  and  $15.998 \pm 0.003$  eV respectively.

Fig. 3 shows the HeI spectrum at a stagnation pressure of 150 torr and a nozzle temperature of 80 K. The spectrum shows well resolved vibrational fine structures on both  $A^2\Sigma^+_{1/2u}$  and  $B^2\Pi_{3/2g}$  states. As many as 14 vibrational excitations are resolved in the ground ionic state. The observed IEs correlate very well with those calculated using the expression,  $E_v = E_0 + \omega_e(v + 1/2) - \omega_e x_e(v + 1/2)^2$  where the values of  $E_0$ ,  $\omega_e$  and  $\omega_e x_e$  are from the work of Morioka et al<sup>12</sup>. The observed vibrational progression is assigned to be due to  $v = 32 - 45$ . The observed ionization energies with the assignments are listed in Table I. Using a Franck-Condon analysis of the photoelectron spectrum, Dehmer and Dehmer<sup>4</sup> estimate that the values of  $\omega_e$  and  $\omega_e x_e$  to be  $308.9$  and  $1.658 \text{ cm}^{-1}$  respectively. The dissociation energy ( $D_e$ ) estimated is  $1.361$  eV, close to the value obtained by Dehmer and Dehmer<sup>4</sup> ( $1.349$  eV). An analysis of the ionization energies of the  $B^2\Pi_{3/2g}$  state gives the values of  $\omega_e$  and  $\omega_e x_e$  to be  $201.6$  and  $12.12 \text{ cm}^{-1}$ , respectively with an adiabatic IE of  $15.632$  eV. This gives a dissociation energy of  $0.104$  eV. The observed excitations are assigned to the  $v = 2 - 4$  vibrational levels of the  $B^2\Pi_{3/2g}$  state.

Dissociation energies ( $D_0$ ) of the ionic states can be estimated from the relation,  $D_0(\text{Rg}_2^+) \geq \text{IE}(\text{Rg}) + D_0(\text{Rg}_2) - \text{IE}(\text{Rg}_2)$ , where  $\text{IE}(\text{Rg}_2)$  refers to the adiabatic IE of the state in question. The dissociation energy of the newly observed  $\text{C}^2\Pi_{3/2}$  state is  $0.061 \pm 0.003$  eV. The  $\text{D}^2\Sigma^+_{1/2g}$  state is found to be weakly attractive, contrary to the theoretical expectations. The dissociation energies of the other states reported here are close to the values in the literature.

The photoelectron spectrum is essentially similar to that reported by Dehmer and Dehmer.<sup>4</sup> There are significant improvements, however. Firstly, we have resolved all the ionic states, including the  $\text{C}^2\Pi_{3/2u}$  state. Secondly, vibrational fine structures on two of the ionic states have been resolved. Although a vibrational progression has been observed on the  $^2\Sigma^+_{1/2u}$  state before, there has been no report hitherto on the vibrational structure of the  $\text{B}^2\Pi_{3/2g}$  state. This is the first reported observation of the vibrational structure of a van der Waals molecule by HeI photoelectron spectroscopy. The ionization energies, dissociation energies and assignments are listed in Table II. Spectroscopic constants of the ground and the first excited ionic states are presented in Table III.

### B. Kr<sub>2</sub>

The photoelectron spectrum of a mixture of Kr and  $\text{Kr}_2$  at a stagnation pressure of 800 torr and at a nozzle temperature of 300 K is shown in Fig. 4. The photoelectron spectrum of  $\text{Kr}_2$  clearly shows all the ionic states of the dimer. The features marked 1, 2, 3, 4, 5 and 6 are assigned to the dimer based on the considerations discussed above. The dimer features were seen even at a stagnation pressure of 300 torr. The intensity of the features increased as the stagnation pressure increased, up to a value of 1200 torr. The states marked 1, 2 and 5 have been observed before by Dehmer and Dehmer<sup>4</sup>. The spectrum is very similar to that of  $\text{Ar}_2$ , with a significant difference being the observation of the  $\text{C}^2\Pi_{3/2u}$  state at the higher IE side of the atomic  $^2P_{3/2}$  line,

implying that the state may be repulsive unlike  $\text{Ar}_2$ , in which the state is weakly bound. The appearance energy for  $\text{Kr}_2$  for the ground state ion is  $13.561 \pm 0.003$  eV. Although the value of  $\omega_e$  is in the range of instrumental resolution<sup>4</sup>, no vibrational excitations were observed. Table IV summarizes the experimental results.

### C. $\text{Xe}_2$

Fig. 5 shows the photoelectron spectrum of a mixture of Xe and  $\text{Xe}_2$ . The spectrum in general agrees with that reported by Dehmer and Dehmer<sup>5</sup>. However, it can be seen that in addition to the states observed by them, the spectrum also shows a feature due to the  $\text{D}^2\Sigma^+_{1/2g}$  state. This state has been observed in the resonantly enhanced photoelectron spectrum of Dehmer et al<sup>10</sup> also. Our estimate of the adiabatic IE of this state is  $13.434 \pm 0.003$  eV, close to the value of 13.42 eV of Dehmer et al<sup>10</sup>. A semi-log plot of the experimental data in the  $^2\text{P}_{3/2}$  region (Fig. 6) shows the  $\text{C}^2\Pi_{3/2u}$  state. Our best estimate adiabatic IE for this state is  $12.053 \pm 0.003$  eV, giving a dissociation energy ( $D_0$ ) of  $0.100 \pm 0.003$  eV. Table V summarizes the experimental results.

The dissociation energies of the rare gas dimers show a general trend. The ground state ion becomes less attractive from  $\text{Ar}_2$  to  $\text{Xe}_2$ . The  $\text{B}^2\Pi_{3/2g}$  state follows a reverse order, more attractive in  $\text{Xe}_2$  and less in  $\text{Ar}_2$ . The same trend is exhibited by the  $\text{C}^2\Pi_{1/2u}$  state as well. The  $\text{D}^2\Sigma^+_{1/2g}$  state of  $\text{Ar}_2$  is only very weakly bound. The dissociation energies of this state for  $\text{Kr}_2$  and  $\text{Xe}_2$  are higher. The second and third excited states ( $\text{C}^2\Pi_{3/2u}$  and  $\text{B}^2\Pi_{1/2g}$ ) are either weakly attractive or weakly repulsive.

In conclusion, the present study gives the ionization energies of all the ionic states of the rare gas dimers in the HeI energy region. Ionization energies of the previously known states are obtained at a much better accuracy. Many of the vibrational structures of the ground and the first ionic states of  $\text{Ar}_2$  are resolved and are

assigned as due to excitations to the higher vibrational levels of the respective states. The study has provided accurate values of the spectroscopic constants of these states. The fifth excited ionic state ( $D^2\Sigma^+_{1/2g}$ ) is weakly bound in all the dimers.

## ACKNOWLEDGMENTS

This work was supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Chemical Science Division of the U.S. Department of Energy under contract No. DE-AC03-76SF00098. We acknowledge the earlier studies of Drs. James Pollard and Dennis Trevor of this group on the vibrational structure of the ground state of  $Ar_2^+$ ; these unpublished earlier spectra and their preliminary analysis, are consistent with our present results. We thank them for reading the manuscript and for critical suggestions. One of the authors (T.P) acknowledges the receipt of an Indo-US science and technology fellowship and thanks the Department of Science and Technology, Government of India and the USAID for financial support. He also thanks all members of the Dave Shirley group for making his stay in the United States enjoyable.

---

<sup>1</sup>Indo-US science and technology fellow, on leave from : CSIR Centre of Excellence in Chemistry, Indian Institute of Science, Bangalore 560 012, INDIA.

<sup>2</sup> Present address: The Pennsylvania State University, 114 Kern Graduate Building, University Park, PA 16802 USA.



**References**

- <sup>1</sup> R. S. Mulliken, *J. Chem. Phys.*, **52**, 5170 (1970).
- <sup>2</sup> W. R. Wadt, *J. Chem. Phys.*, **68**, 402 (1978).
- <sup>3</sup> H. H. Michels, R. H. Hobbs and L. A. Wright, *J. Chem. Phys.*, **69**, 5151 (1978).
- <sup>4</sup> P. M. Dehmer and J. L. Dehmer, *J. Chem. Phys.*, **69**, 125 (1978).
- <sup>5</sup> P. M. Dehmer and J. L. Dehmer, *J. Chem. Phys.*, **68**, 3462 (1978).
- <sup>6</sup> J. T. Moseley, R. P. Saxson, B. A. Huber, P. C. Cosby, R. Abouaf and M. Tadjeddine, *J. Chem. Phys.*, **67**, 1659 (1977).
- <sup>7</sup> C. Y. Ng, D. J. Trevor, B. H. Mahan and Y. T. Lee, *J. Chem. Phys.*, **65**, 4327 (1976).
- <sup>8</sup> C. Y. Ng, D. J. Trevor, B. H. Mahan and Y. T. Lee., *J. Chem. Phys.*, **66**, 446 (1977).
- <sup>9</sup> R. Abouaf, B. A. Huber, P. C. Cosby, R. P. Saxson and J. T. Moseley, *J. Chem. Phys.*, **68**, 2406 (1978).
- <sup>10</sup> P. M. Dehmer, S. T. Pratt and J. L. Dehmer, *J. Phys. Chem.*, **91**, 2593 (1987).
- <sup>11</sup> R. J. LeRoy, *J. Chem. Phys.*, **57**, 573 (1972).
- <sup>12</sup> Y. Morioka, M. Ogawa, T. Matsumoto, K. Ito, K. Tanaka and T. Hayaishi, *J. Phys. Ato. Mol. Opt. Phys.*, **24**, 791 (1991).
- <sup>13</sup> R. G. Tonkyn and M. G. White, *J. Chem. Phys.*, **95**, 5582 (1991).
- <sup>14</sup> J. E. Pollard, D. J. Trevor, Y. T. Lee and D. A. Shirley, *Rev. Sci. Instrum.*, **52**, 1837 (1981).
- <sup>15</sup> J. A. Barker, R. O. Watts, J. K. Lee, T. P. Schafer and Y. T. Lee, *J. Chem. Phys.*, **61**, 3081 (1974).
- <sup>16</sup> P. M. Dehmer and S. T. Pratt., *J. Chem. Phys.*, **76**, 843 (1982).
- <sup>17</sup> F. Carnovale, J. B. Peel, R. G. Rothwell, J. Valdorf and P. J. Kunz, *J. Chem. Phys.*, **90**, 1452 (1989).
- <sup>18</sup> O. F. Hagena, *Z. Phys.*, **D4**, 291 (1987).

<sup>19</sup> J. Wormer, V. Guzielski, J. Stapelfeldt, G. Zimmerer and T. Moller, *Physica Scripta*, **41**, 490 (1990).

<sup>20</sup> T. L. Gilbert and A. C. Wahl, *J. Chem. Phys.*, **55**, 5247 (1977).

<sup>21</sup> W. J. Stevens, M. Gardner, A. Kara and P. Julienne, *J. Chem. Phys.*, **67** (1977) 2860.

TABLE I. Ionization potentials of the  $A^2\Sigma^+_{1/2u}$  state observed in the HeI photoelectron spectrum.

Observed <sup>a</sup>	IE (eV)	Calculated <sup>b</sup>	$\nu$	$\Delta G(\nu + 1/2)$ (cm <sup>-1</sup> )
15.396(1)		15.403	32	-
15.419(1)		15.424	33	189
15.440(1)		15.444	34	165
15.461(1)		15.463	35	166
15.481(1)		15.482	36	166
15.498(1)		15.500	37	137
15.514 (1)		15.518	38	132
15.531(3)		15.535	39	132
15.548(3)		15.552	40	137
15.564(1)		15.568	41	132
15.579(1)		15.584	42	120
15.592(1)		15.599	43	108
15.603(1)		15.613	44	84
15.612(1)		15.627	45	77

<sup>a</sup> The numbers in the parentheses are the uncertainties in the last digit.

<sup>b</sup> Using the relation,  $E_\nu = E_0 + \omega_e(\nu + 1/2) - \omega_e x_e(\nu + 1/2)^2$ . The values of  $E_0$ ,  $\omega_e$  and  $\omega_e x_e$  are from ref. 12. The difference between the observed and calculated IEs is large for the last three IEs, possibly because Morse function is a poor approximation in describing the  $A^2\Sigma^+_{1/2u}$  state near the dissociation limit.

TABLE II. Ionization potentials and dissociation energies<sup>a</sup> of the ionic states of Ar<sub>2</sub> observed in the HeI energy region.

Peak State	Adiabatic IE (eV)		Vertical IE (eV)		Dissociation Energy (eV)		
	Present	Ref. 4	Present	Ref. 4	Present	Ref. 4	Other
1 A <sup>2</sup> Σ <sup>+</sup> <sub>1/2u</sub>	.b	.b	15.548±0.003	15.55±0.02			1.19 <sup>c</sup> 1.30 <sup>d</sup> 1.33 <sup>e</sup> 1.25 <sup>f</sup> 1.20 <sup>g</sup> 1.27 <sup>h</sup>
2 B <sup>2</sup> Π <sub>3/2g</sub>	15.630±0.003	15.63±0.02	15.689±0.001	15.67±0.02	0.139±0.003	0.14±0.02	0.10 <sup>c</sup> 0.16 <sup>d</sup>
3 C <sup>2</sup> Π <sub>3/2u</sub>	15.708±0.003		15.734±0.003		0.061±0.003		0.01 <sup>c</sup> 0.03 <sup>d</sup>
4 B <sup>2</sup> Π <sub>1/2g</sub>			15.829±0.001		repulsive		repulsive <sup>c</sup> 0.05 <sup>d</sup>
5 C <sup>2</sup> Π <sub>1/2u</sub>	15.847±0.003	15.84±0.015	15.871±0.001	15.87±0.015	0.100±0.003	0.100±0.015	0.04 <sup>c</sup> 0.08 <sup>d</sup>
6 D <sup>2</sup> Σ <sup>+</sup> <sub>1/2g</sub>	15.934±0.003		15.998±0.003	15.99±0.03	0.013±0.003	repulsive	repulsive <sup>c,d</sup>

## Table II (cont.)

- <sup>a</sup> The experimental values correspond to  $D_0$  not  $D_e$ .
- <sup>b</sup> The adiabatic IE is not observed due to poor Franck-Condon factors.
- <sup>c</sup> From Wadt (ref. 2), the value corresponds to  $D_e$  not  $D_0$ .
- <sup>d</sup> From Michels et al (ref. 3).
- <sup>e</sup> From Moseley et al (ref. 6).
- <sup>f</sup> From Gilbert and Wahl (ref. 20).
- <sup>g</sup> From Stevens et al (ref. 21).
- <sup>h</sup> From the photoionization studies of Dehmer and Pratt (ref. 16) which gives a value of 14.49 eV for the adiabatic IE.

TABLE III. Spectroscopic parameters of  $\text{Ar}_2^+$  for the  $A^2\Sigma^+_{1/2u}$  and the  $B^2\Pi_{3/2g}$  states.

State	$D_e$ (eV) <sup>a</sup>	$\omega_e$ (cm <sup>-1</sup> ) <sup>b</sup>	$\omega_e x_e$ (cm <sup>-1</sup> ) <sup>b</sup>
$A^2\Sigma^+_{1/2u}$	1.361 <sup>c,d</sup>	310.8 <sup>c,d</sup>	2.2 <sup>c,d</sup>
$B^2\Pi_{3/2g}$	0.104	201.6	12.12

<sup>a</sup> Using the relation,  $D_e = \omega_e^2/4\omega_e x_e$

<sup>b</sup> The values of  $\omega_e$  and  $\omega_e x_e$  are obtained by least squares fitting the observed ionization energies of the respective states to the relation,  $E_v = E_0 + \omega_e (v + 1/2) - \omega_e x_e (v + 1/2)^2$ .

<sup>c</sup> From Morioka et al (ref. 12).

<sup>d</sup> Dehmer and Dehmer (ref. 4 ) give a value of 1.349, 308.9 and 1.658 for  $D_e$ ,  $\omega_e$  and  $\omega_e x_e$  respectively on the basis of a Frack-Condon analysis of the observed photoelectron peak.

TABLE IV. Ionization potentials and dissociation energies<sup>a</sup> of the ionic states of Kr<sub>2</sub> observed in the HeI energy region.

Peak	State	Adiabatic IE (eV)		Vertical IE (eV)		Dissociation Energy (eV)		
		Present	Ref. 4	Present	Ref. 4	Present	Ref. 4	Other
1	A <sup>2</sup> Σ <sup>+</sup> <sub>1/2u</sub>	.b	.b	13.773±0.003	13.76±0.02			1.05 <sup>c</sup> 1.18 <sup>d</sup> 1.176 <sup>e</sup> 1.15 <sup>f</sup>
2	B <sup>2</sup> Π <sub>3/2g</sub>	13.831±0.003	13.85±0.0015	13.907±0.001	13.90±0.015	0.184±0.003	0.16±0.015	0.13 <sup>c</sup> 0.16 <sup>d</sup>
3	C <sup>2</sup> Π <sub>3/2u</sub>			14.055±0.003		repulsive		0.02 <sup>c</sup> 0.07 <sup>d</sup>
4	B <sup>2</sup> Π <sub>1/2g</sub>			14.117±0.003		repulsive		repulsive <sup>c,d</sup>
5	C <sup>2</sup> Π <sub>1/2u</sub>	14.530±0.003	14.54±0.015	14.566±0.001	15.57±0.015	0.151±0.003	0.14±0.015	0.17 <sup>c</sup> 0.10 <sup>d</sup>
6	D <sup>2</sup> Σ <sup>+</sup> <sub>1/2g</sub>	14.653±0.003		14.728±0.003		0.028±0.003		repulsive <sup>c,d</sup>

<sup>a</sup> The experimental values correspond to D<sub>0</sub> not D<sub>e</sub>.

<sup>b</sup> The adiabatic IE is not observed due to poor Franck-Condon factors.

<sup>c</sup> From Wadt (ref. 2), the value corresponds to D<sub>e</sub> not D<sub>0</sub>.

<sup>d</sup> From Michels et al (ref. 3). <sup>e</sup> From Abouaf et al (ref. 9). <sup>f</sup> From Ng et al (ref. 8).

TABLE V. Ionization potentials and dissociation energies<sup>a</sup> of the ionic states of Xe<sub>2</sub> observed in the HeI energy region.

Peak State	Adiabatic IE (eV)		Vertical IE (eV)		Dissociation Energy (eV)		
	Present	Ref. 4	Present	Ref. 4	Present	Ref. 4	Other
1 A <sup>2</sup> Σ <sup>+</sup> <sub>1/2u</sub>	.b	.b	11.845±0.003	11.85±0.015			0.79 <sup>c</sup> 1.06 <sup>d</sup> 1.03 <sup>e</sup> 0.75 <sup>f</sup>
2 B <sup>2</sup> Π <sub>3/2g</sub>	11.947±0.003	11.97±0.0015	12.021±0.001	12.02±0.015	0.207±0.003	0.185±0.015	0.12 <sup>c</sup> 0.15 <sup>d</sup> 0.21 <sup>f</sup>
3 C <sup>2</sup> Π <sub>3/2u</sub>	12.053±0.003		12.089±0.001 <sup>g</sup>		0.100±0.003		0.03 <sup>c</sup> 0.04 <sup>d</sup> 0.05 <sup>f</sup>
4 B <sup>2</sup> Π <sub>1/2g</sub>	12.139±0.003		12.198±0.001	12.21±0.015	0.014±0.003	repulsive	repulsive <sup>c,d</sup>
5 C <sup>2</sup> Π <sub>1/2u</sub>	13.263±0.003	13.27±0.015	13.316±0.001	13.31±0.015	0.196±0.003	0.19±0.015	0.12 <sup>c</sup> 0.21 <sup>d</sup> 0.17 <sup>f</sup>
6 D <sup>2</sup> Σ <sup>+</sup> <sub>1/2g</sub>	13.434±0.003		13.518±0.003		0.025±0.003		repulsive <sup>c,d</sup> 0.04 <sup>f</sup>



**Table V (Cont.)**

<sup>a</sup> The experimental values correspond to  $D_0$  not  $D_e$ .

<sup>b</sup> The adiabatic IE is not observed due to poor Franck-Condon factors.

<sup>c</sup> From Wadt (ref. 2), the value corresponds to  $D_e$  not  $D_0$ .

<sup>d</sup> From Michels et al (ref. 3).

<sup>e</sup> From Ng et al (ref. 7).

<sup>f</sup> From Dehmer et al (ref. 10).

<sup>g</sup> Dehmer et al (Ref. 10) give a value of 12.01 eV for the vertical IE.

### FIGURE CAPTIONS

**FIG. 1.** He I photoelectron spectrum of a mixture of Ar and Ar<sub>2</sub> at a stagnation pressure of 280 torr and at a nozzle temperature of 80 K with an instrumental resolution of 13 meV FWHM. The spectrum has been truncated to enhance the dimer features which are 100 times weaker than the atomic  $2P_{3/2}$  peak. The dimer features are labeled 1, 2, 4, 5 and 6. Weak features due to the impurity lines in the undispersed light source are also seen. The vibrational fine structures on features 1 and 2 are marked with vertical lines. A gradual rise in the intensity above the background is seen beginning  $15.138 \pm 0.003$  eV which is identified as the appearance energy of the dimer.

**FIG. 2.** A semi-log plot of the data presented in Fig. 1 near the dimer features. The feature labeled 3 corresponds to the  $C^2\Pi_{3/2g}$  state.

**FIG. 3.** HeI photoelectron spectrum of a mixture of Ar and Ar<sub>2</sub> at a stagnation pressure of 150 torr and at a nozzle temperature of 80 K. The vibrational structures of the  $A^2\Sigma^+_{1/2u}$  and the  $B^2\Pi_{3/2g}$  states are marked with vertical lines. For the  $A^2\Sigma^+_{1/2u}$  state 14 vibrational states are visible. Only three states are resolved for the  $B^2\Pi_{3/2g}$  state.

**FIG. 4.** The HeI photoelectron spectrum of a mixture of Kr and Kr<sub>2</sub> at a nozzle stagnation pressure of 800 torr and at a nozzle temperature of 300 K. The labeling of the features is as in Fig. 1.

**FIG. 5.** The HeI photoelectron spectrum of a mixture of Xe and Xe<sub>2</sub> at a nozzle stagnation pressure of 820 torr and at a nozzle temperature of 300 K. The labels have the same meaning as in Fig. 1.

**FIG. 6.** A semi-log plot of the data presented in Fig. 5 in the Xe  $2P_{3/2}$  region. The feature marked 3 is due to the  $C^2\Pi_{3/2u}$  state. Wiggles in the spectrum in regions of low intensity are due to data averaging.




FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6



LAWRENCE BERKELEY LABORATORY  
UNIVERSITY OF CALIFORNIA  
TECHNICAL INFORMATION DEPARTMENT  
BERKELEY, CALIFORNIA 94720